

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number
WO 01/96607 A2

- (51) International Patent Classification⁷: C12Q 1/68
- (21) International Application Number: PCT/US01/19249
- (22) International Filing Date: 13 June 2001 (13.06.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/211,356 13 June 2000 (13.06.2000) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/211,356 (CIP)
Filed on 13 June 2000 (13.06.2000)
- (71) Applicant (for all designated States except US): THE TRUSTEES OF BOSTON UNIVERSITY [US/US]; 108 Bay State Road, Boston, MA 02215 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): CANTOR, Charles, R. [US/US]; 526 Stratford Court, Unit E, Del Mar, CA 92014 (US). SIDDIQI, Fouad, A. [US/US]; Apt. #3, 113 Bay State Road, Boston, MA 02215 (US).
- (74) Agents: SEIDMAN, Stephanie, L. et al.; Heller Ehrman White & McAuliffe LLP, Suite 700, 4250 Executive Square, La Jolla, CA 92037 (US).
- (81) Designated States (national): AE, AG, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/96607 A2

(54) Title: USE OF NUCLEOTIDE ANALOGS IN THE ANALYSIS OF OLIGONUCLEOTIDE MIXTURES AND IN HIGHLY MULTIPLEXED NUCLEIC ACID SEQUENCING

(57) Abstract: Methods and kits that use nucleotide analogs to confer increased accuracy and improved resolution in the analysis and sequencing of oligonucleotide mixtures are provided.

BEST AVAILABLE COPY

USE OF NUCLEOTIDE ANALOGS IN THE ANALYSIS OF OLIGONUCLEOTIDE MIXTURES AND IN HIGHLY MULTIPLEXED NUCLEIC ACID SEQUENCING

Subject matter described herein was developed under NSF Grant No. Ger-9452651. The Government can have certain rights therein.

5 RELATED APPLICATIONS

For U.S. purposes for priority is claimed under 35 U.S.C. §119(e) to U.S. provisional application Serial No. 60/211,356, filed June 13, 2000, to Charles R. Cantor and Fouad A. Siddiqi, entitled USE OF NUCLEOTIDE ANALOGS IN THE ANALYSIS OF OLIGONUCLEOTIDE MIXTURES AND IN HIGHLY MULTIPLEXED
10 NUCLEIC ACID SEQUENCING." For international purposes benefit of priority is claimed thereto. The subject matter of U.S. provisional application Serial No. 60/211,356 is incorporated by reference in its entirety.

FIELD OF THE INVENTION

This invention relates to methods, particularly mass spectrometric
15 methods, for the analysis and sequencing of nucleic acid molecules.

DESCRIPTION OF THE BACKGROUND

Since the recognition of nucleic acid as the carrier of the genetic code, a great deal of interest has centered around determining the sequence of that code in the many forms in which it occurs. Two studies made the process of nucleic
20 acid sequencing, at least with DNA, a common and relatively rapid procedure practiced in most laboratories. The first describes a process whereby terminally labeled DNA molecules are chemically cleaved at single base repetitions (A.M. Maxam and W. Gilbert, *Proc. Natl. Acad. Sci. USA* 74:560-64, 1977). Each base position in the nucleic acid sequence is then determined from the molecular
25 weights of fragments produced by partial cleavage. Individual reactions were devised to cleave preferentially at guanine, at adenine, at cytosine and thymine, and at cytosine alone. When the products of these four reactions are resolved by molecular weight, using, for example, polyacrylamide gel electrophoresis, DNA sequences can be read from the pattern of fragments on the resolved gel.

30 In another method DNA is sequenced using a variation of the plus-minus method (Sanger *et al.* (1977) *Proc. Natl. Acad. Sci. USA* 74:5463-67, 1977). This procedure takes advantage of the chain terminating ability of

-2-

dideoxynucleoside triphosphates (ddNTPs) and the ability of DNA polymerase to incorporate ddNTPs with nearly equal fidelity as the natural substrate of DNA polymerase, deoxynucleoside triphosphates (dNTPs). Briefly, a primer, usually an oligonucleotide, and a template DNA are incubated in the presence of a
5 useful concentration of all four dNTPs plus a limited amount of a single ddNTP. The DNA polymerase occasionally incorporates a dideoxynucleotide that terminates chain extension. Because the dideoxynucleotide has no 3'-hydroxyl, the initiation point for the polymerase enzyme is lost. Polymerization produces a mixture of fragments of varied sizes, all having identical 3' termini. Fractionation
10 of the mixture by, for example, polyacrylamide gel electrophoresis, produces a pattern that indicates the presence and position of each base in the nucleic acid. Reactions with each of the four ddNTPs permits the nucleic acid sequence to be read from a resolved gel.

These procedures are cumbersome and are limited to sequencing DNA.
15 In addition, with conventional procedures, individual sequences are separated by, for example, electrophoresis using capillary or slab gels, which slow. Mass spectrometry has been adapted and used for sequencing and detection of nucleic acid molecules (see, *e.g.*, U.S. Patent Nos. (6,194,144; 6,225,450; 5,691,141; 5,547,835; 6,238,871; 5,605,798; 6,043,031; 6,197,498; 6,235,478;
20 6,221,601; 6,221,605). In particular, Matrix-Assisted Laser Desorption/Ionization (MALDI) and ElectroSpray Ionization (ESI), which allow intact ionization, detection and exact mass determination of large molecules, *i.e.* well exceeding 300 kDa in mass have been used for sequencing of nucleic acid molecules.

25 A further refinement in mass spectrometric analysis of high molecular weight molecules was the development of time of flight mass spectrometry (TOF-MS) with matrix-assisted laser desorption ionization (MALDI). This process involves placing the sample into a matrix that contains molecules that assist in the desorption process by absorbing energy at the frequency used to desorb the
30 sample. Time of flight analysis uses the travel time or flight time of the various ionic species as an accurate indicator of molecular mass. Due to its speed and high resolution, time-of-flight mass spectrometry is well-suited to the task of

-3-

short-range, *i.e.*, less than 30 base sequencing of nucleic acids. Since each of the four naturally occurring nucleotide bases dC, dT, dA and dG, also referred to herein as C, T, A and G, in DNA has a different molecular weight,

$$M_C = 289.2$$

5 $M_T = 304.2$

$$M_A = 313.2$$

$$M_G = 329.2,$$

where M_C , M_T , M_A , M_G are average molecular weights in daltons of the nucleotide bases deoxycytidine, thymidine, deoxyadenosine, and
10 deoxyguanosine, respectively, it is possible to read an entire sequence in a single mass spectrum. If a single spectrum is used to analyze the products of a conventional Sanger sequencing reaction, where chain termination is achieved at every base position by the incorporation of dideoxynucleotides, a base sequence
15 can be determined by calculation of the mass differences between adjacent peaks. In addition, the method can be used to determine the masses, lengths and base compositions of mixtures of oligonucleotides and to detect target oligonucleotides based upon molecular weight.

MALDI-TOF mass spectrometry for sequencing DNA using mass modification (see, *e.g.*, U.S. Patent Nos. 5,547,835, 6,194,144; 6,225,450;
20 5,691,141 and 6,238,871) to increase mass resolution is available. The methods employ conventional Sanger sequencing reactions with each of the four dideoxynucleotides. In addition, for example for multiplexing, two of the four natural bases are replaced; dG is substituted with 7-deaza-dG and dA with 7-deaza-dA.

25 A variety of techniques and combinations thereof have been directed to improving the level of accuracy in determining the nucleotide compositions of mixtures of oligonucleotides using mass spectrometry, and many of these methods employ nucleotide analogs. For example, Muddiman *et al.* (*Anal. Chem.*, 69(8): 1543-1549, 1997) discusses an algorithm for the unique
30 definition of the base composition of PCR-amplified products, especially longer (> 100bp) oligonucleotides. The algorithm places a constraint on the otherwise large number of possible base compositions for long oligonucleotides by taking

-4-

into account only those masses (measured by electrospray ionization mass spectrometry) that are consistent with that of their denatured complementary strands, assuming Watson-Crick base-pairing. In addition, the algorithm imposes the constraint of known primer compositions, since the primer sequences are known, and this constraint becomes especially significant with shorter PCR products whose mass of "unknown" sequence relative to that of the primer mass is small. Muddiman *et al.* also discusses invoking additional measurements for defining the base composition with even greater accuracy. These include the possibility of post-modifying the PCR product using *e.g.*, dimethyl sulfate to selectively methylate every "G" in the PCR product, or using a modified base during PCR amplification, conducting mass measurements on the modified oligonucleotides, and comparing the mass measurements with those of the unmodified complementary strands.

Chen *et al.* (*Anal. Chem.*, 71(15): 3118-3125, 1999) reports a method that combines stable isotope $^{13}\text{C}/^{15}\text{N}$ labelling of PCR products with analysis of the mass shifts by MALDI-TOF mass spectrometry. The mass shift due to labelling of a single type of nucleotide (i.e., A, T, G or C) reveals the number of that type of nucleotide in a given fragment. While the method is useful in the measurement and comparison of nucleotide compositions of homologous sequences for sequence validation and in scoring polymorphisms, tedious repetitive sequencing reactions (using the four different labelled nucleotides) and mass spectrometric measurements are required.

Hence there is a need in the art for methods that (i) unambiguously assign nucleotides in a sequence, and, (ii) resolve large numbers of oligonucleotides that have the same length, different base compositions, and nearly equal (i.e., less than or equal to about 1 dalton difference) molecular weights. Therefore it is an object herein to provide methods that solve such problems

SUMMARY OF THE INVENTION

Provided herein are methods for sequencing and detecting nucleic acids using techniques, such as mass spectrometry and gel electrophoresis, that are based upon molecular mass. The methods use deoxynucleotide analogs, modified nucleotide terminators and/or mass-labeled primers in one or more

-5-

reactions for sequencing or detection protocols that involve primer extension, and analyze these products from more than one oligonucleotide on, for example, a single mass spectrum. This provides a means for accurate detection and/or sequencing of a an oligonucleotide and is particularly advantageous for detecting
5 or sequencing a plurality target nucleic acid molecules in a single reaction using any technique that distinguishes products based upon molecular weight. The methods herein are particularly adapted for mass spectrometric analyses.

For example, a sequencing method provided herein uses deoxynucleotide analogs, modified nucleotide terminators and/or mass-labeled primers in one or
10 more Sanger sequencing reactions, and analyzes these products from more than one oligonucleotide on a single mass spectrum. In particular, a plurality of primers can be used to simultaneously sequence a plurality of nucleic acid molecules or portions of the same molecule. By extending the primers with mass-matched nucleotides, the resulting products mass shifts that are
15 periodically related to the size of the original primer.

As a result, the sequence of any given oligonucleotide can be determined with a high level of accuracy, and also mixtures of a number of sequences can be multiplexed in a single mass spectrum. The limit on the number of oligonucleotides that can be sequenced simultaneously is governed by the base
20 periodicity, the maximum mass shift, and the resolving power of analytical tool, such as the mass spectrometer. The base periodicity and maximum mass shift can be carefully engineered for optimal resolution and accuracy, depending on the number of sequences to be simultaneously analyzed, and the information desired; as many sequences as desired can be sequenced simultaneously
25 especially in the detection and scoring of single nucleotide polymorphisms, insertions, deletions and other mutations.

In another embodiment, a target nucleic acid molecule is sequenced using mass-matched nucleotides and chain terminating nucleotides. For example, a primer is annealed to a target nucleic acid, the primer is extended in
30 the presence of chain-terminating nucleotides and mass-matched nucleotides to produce extension products, the masses of the extension products follow a periodic distribution that is determined by the mass of the mass-matched

-6-

nucleotides, and the sequence of the target nucleic acid is determined from the mass shift of each extension product from its corresponding periodic reference mass by virtue of incorporation of the chain terminator. The mass-matched nucleotides all have identical masses, and each chain terminating nucleotide has a distinct mass that differs from that of the other chain terminating nucleotides. This results in unique predetermined values of mass shift corresponding to each chain terminating nucleotide and based upon the original primer.

This method is adaptable for any sequencing method or detection method that relies upon or includes chain extension. These methods include, but are not limited to, sequencing methods based upon Sanger sequencing, and detection methods, such as primer oligo base extension (PROBE) (see, *e.g.*, U.S. application Serial No. 6,043,031; allowed U.S. application Serial No. 09/287,679; and 6,235,478), that rely include a step of chain extension. Also, contemplated are methods, such as haplotyping methods, in which two mutations in the same gene are detection are provided. A detector (primer) oligonucleotide is to the hybridized to the first mutation and the primer is extended with mass-matched nucleotides and appropriately selected chain terminator(s) to detect the second mutation.

In other embodiments, a plurality of target nucleic acids can be multiplexed in a single reaction measurement by annealing each target nucleic acid to a primer of distinct molecular weight each primer is then extended with mass-matched nucleotides and chain terminators in formats that depend upon whether detection or sequencing is desired. These methods are particularly useful for methods of detection in which a primer is hybridized to a plurality of target nucleic acid molecules, such as immobilized nucleic acid molecules, hybrids separated from unhybridized nucleic acids and the detectors detected. Such methods include PROBE, in which case the extension reaction is performed in the presence chain terminators and mass matched deoxynucleotides.

The primers of distinct molecular weight can be selected to differ in molecular weight by a value that is greater than the maximum mass shift, *i.e.*, the difference in molecular weight between the heaviest and the lightest nucleotide terminators in chain extension reactions. The difference in molecular

-7-

weight between the primers for a plurality of target nucleic acids can be selected to be least 20 daltons greater than the maximum mass shift to account for the finite band width of the peaks.

The number of molecules that can be multiplexed is governed by the
5 periodicity, the maximum mass shift, and the resolving power of the sequence detection instrument. In some embodiments, about 7 to about 25 or more molecules can be multiplexed. For scoring single nucleotide polymorphisms, only a single nucleotide terminator is required (depending on the base identity of the single nucleotide polymorphism). In this case, the maximum mass shift required
10 is identically zero, so that larger numbers of molecules, greater than 25, 35, 50 and more, can be multiplexed, depending on the resolving power of the sequencing format, and for mass spectrometry the instrument. Depending on the amount of sequence information desired, one, two or three rather than four types of nucleotide terminators (corresponding to each of the four nucleic acid
15 bases) can be used.

In other embodiments, the mass shift is obtained using pair-matched nucleotides, *i.e.*, the mass of each nucleotide base-pair is selected so that the masses of all pairs are identical. In one embodiment thereof, the following steps are performed: (i) the target nucleic acid is copied or amplified by a method such
20 as PCR in the presence of the pair-matched nucleotide set prior to the sequencing or detection reaction; (ii) the target nucleic acid is denatured, and a partially duplex hairpin primer is annealed and ligated to the single-stranded template; (iii) the primer is extended in the presence of chain terminating nucleotides and pair-matched nucleotides to produce extension products, where
25 the masses of the extension products follow a periodic distribution that is determined by the mass of the pair-matched nucleotide set, and, (iv) the target nucleic acid is detected by virtue of its molecular weight or its sequence is determined from the mass shift of each extension product from its corresponding periodic reference mass.

-8-

In another embodiment, the mass of each terminating base pair is unique and resolvable, so that the mass shifts corresponding to each terminating base pair are unique. The nucleotide terminators are optionally mass-matched or can be of distinct masses as long as distinct values of mass shift are obtained for each terminating base pair.

In another embodiment, the extension products are treated to produce blunt-ended double-stranded extension products by methods known to those of skill in the art, such as the use of single-strand specific nucleases. In an aspect of this embodiment, a plurality of target nucleic acids can be multiplexed in a single reaction by annealing each target nucleic acid to a primer of distinct molecular weight. The primers can be selected to differ in molecular weight by a value that is greater than the maximum mass shift, *i.e.*, the difference in molecular weight between the heaviest and the lightest nucleotide terminating base pairs. Since double stranded nucleic acid can be analyzed, the effective sequence read is halved relative to the embodiment employing mass-matched nucleotides, but the number of molecules that can be multiplexed is doubled, due to the increase in period (the value of the mass of a base pair, rather than a single mass-matched nucleotide). In exemplary embodiments, about 14 to about 50 sequences are multiplexed. In detection embodiments, about 50 or more molecules can be simultaneously multiplexed since only a single terminating base pair is added in the extension reaction.

In another embodiment, the chain termination reactions are carried out separately using a standard nucleotide terminator, pair-matched nucleotides, and mass-labeled primers, if modified nucleotide terminators which are either mass-matched or provide distinct values of mass shift for each terminating base pair are not available. The reactions are pooled prior to detection or sequence analysis. In one embodiment, the mass-labeled primers can have distinct values of molecular weight that give rise to unique values of mass shift or positional mass difference for each terminating base.

In another method provided herein, a population of nucleic acids having the same length but different base compositions can be resolved by synthesizing the nucleic acids in the presence of a nucleotide analog to produce synthesized

-9-

nucleic acids having incorporated the nucleotide analog, where the nucleotide analog is selected to optimally separate the masses of the population of nucleic acids according to their individual base compositions. For example, the nucleotide analog or analogs are selected to separate the population of nucleic acids according to base composition by greater than 1 dalton. In another embodiment, the nucleotide analog or analogs are selected to separate the population of nucleic acids according to base composition by mass values of about 3 daltons to about 8 daltons, depending on the choice of analog and on the resolving power of the detection instrument. In other embodiments, the nucleotide analog or analogs can be selected to restrict oligonucleotides having the same length to have the same mass, *i.e.*, a peak separation of zero, regardless of differences in base composition, such as in detection methods, where it is desirable to separate populations of oligonucleotides according to their length.

15 Nucleic acid molecules that contain mass-matched nucleotides and/or pair-matched nucleotides are provided.

Also provided are combinations for practicing the methods provided herein. For instance, in one embodiment, the combinations include a set of mass-matched deoxynucleotides. In another embodiment, the combinations a set of pair-matched nucleotides and a set of mass-matched chain terminating nucleotides. In another embodiment, the combination includes a set of pair-matched nucleotides and chain terminating nucleotides which form terminating base pairs of distinctly different molecular weight. In yet another embodiment, the combination includes a set of pair-matched nucleotides and mass-labeled primers. In other embodiments, mass-staggered primers can be added to as optional components.

Kits containing the combinations with optional instructions and/or additional reagents are also provided. The kits contain the reagents as described herein and optionally any other reagents required to perform the reactions. Such reagents and compositions are packaged in standard packaging known to those of skill in the art. Additional vials, containers, pipets, syringes and other

-10-

products for sequencing can also be included. Instructions for performing the reactions can be included.

Also provided herein are methods for optimization of the analysis of base compositions of mixtures of oligonucleotides by mass spectrometry. A single
5 spectrum can be used to resolve a very large number of oligonucleotides having the same length but different molecular weights by incorporating a nucleotide analog into the oligonucleotides in the mixture such that the peaks are no closer than a minimum value called peak separation. The peak separation can be
tailored by careful selection of the nucleotide analog and of a mass spectrometer
10 with the desired resolving power.

The methods herein permit unambiguous and accurate analysis of the sequences or molecular weights of large numbers of oligonucleotides in a single mass spectrum by combining the rapidity of mass spectrometry with the
resolving power of nucleotide analogs which are carefully selected and
15 incorporated into the oligonucleotide mixture according to the desired application.

Other features and advantages will be apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE FIGURES

20 Figure 1 shows that when a single spectrum is used to analyze the products of a conventional Sanger sequencing reaction, where chain termination is achieved at every base position by the incorporation of dideoxynucleotides, the base sequence can be determined by calculation of the mass differences between adjacent peaks (Figures 1a and 1b).

25 Figure 2 shows implementation of forced mass modulation using mass-matched deoxynucleotides. Figure 2a is a simulated mass spectrum showing the products and molecular masses of a reaction carried out with a suitable polymerase in the presence of a mass-matched nucleotide set ("dN") and the four standard dideoxynucleotide terminators. The base periodicity is the mass of
30 dN, or 310 daltons. Figure 2b shows a target second sequence resolved on the same mass spectrum shown in Figure 2a, using a primer heavier by 77 daltons. The peaks corresponding to the reaction products from the first target sequence

-11-

can fall within the the spectrum in Figure 2b, which can never intersect peaks from the second target sequence. This permits unambiguous resolution of both sequences each peak can be uniquely assigned to a nucleotide, a base position, and a target sequence.

5 Figure 3 shows four different sequences resolved in a single spectrum using a set of mass-staggered primers that are separated in mass by integer multiples of 77 daltons (77, 154, and 231 daltons).

10 Figure 4 shows the general implementation of a forced mass modulation method using pair-matched nucleotides, for the analysis of sequencing reaction products as double-stranded structures. The steps in the reaction are as follows:
a) a partially duplex hairpin primer with a 3' overhang and a 5' phosphate group is annealed and ligated to the single stranded target sequence; b) the resulting partially duplex structure is subjected to a sequencing reaction using the pair-matched nucleotide set described above along with the set of mass-matched
15 terminators (ddM); c) products resulting from sequencing reaction b); and, d) the products c) from the sequencing reaction are exposed to a strict single strand-specific nuclease that results in the production of blunt-ended hairpin structures ready for analysis by mass spectrometry.

20 Figure 5 shows the products and molecular masses of the nuclease digestion elucidated in Figure 4d, along with a simulated mass spectrum.

25 Figure 6 shows three sequence variants (Figure 6a) that differ from each other only at a single base position sequenced by a conventional Sanger reaction. Figure 6b is a simulated mass spectrum of all reaction products shown in Figure 6a. Figure 6c is a graph representing the valid sequence permutations that can be elucidated from the mass spectrum shown in Figure 6b. Boxed values are fragment masses, solid arrows show valid sequence branches, dashed arrows represent spurious branches. In practice valid branches are indistinguishable from spurious ones. Figure 6d is a set of sequences consistent with the graph shown in Figure 6c. Spurious sequence reconstructions are shown in lowercase letters, valid ones
30 in uppercase letters.

Figure 7a shows the three related sequence variants from Figure 6 sequenced by Forced Mass Modulation using a single primer and the mass-matched

-12-

nucleotide set from Figure 2 with the standard dideoxy terminators. The positions of the differing bases are shown by solid arrows. Reaction products are shown along with their respective molecular masses. Reaction products of variant #2 whose masses differ from those of variant #1 are marked with by (*). Reaction products of variant #3 whose masses differ from those of variant #1 are marked by (**). Figure 7b is a simulated mass spectrum of all reaction products shown in Figure 7a along with a sequence graph. The shaded regions represent the only valid mass ranges that can assumed by the reaction products from Figure 7a. The base periodicity is 310 daltons. Figure 7c is a consensus sequence derived from the data shown in Figure 7b. Figure 7d is an expansion of the consensus sequence shown in Figure 7c. Spurious reconstructions are shown in lowercase letters, valid ones in uppercase letters. Note that there is only a single spurious reconstruction, as opposed to the eleven errant sequences reconstructed from the Sanger reaction described in Figure 6.

Figure 8 shows the base composition density distributions for the total set of possible 7-base oligonucleotides using three different nucleotide sets. Note that for the set of naturally occurring bases, nearly every base composition has its own distinct mass value, but most of these mass values are spaced only one dalton from each other. Increasing the peak separation also markedly increases the average number of base compositions per observed mass, particularly for those masses in the center of the range.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. All patents, patent applications, Genbank and other sequence repository sequences, and publications referred to herein are incorporated by reference.

As used herein, a biopolymer includes, but is not limited to, nucleic acid, proteins, polysaccharides, lipids and other macromolecules. Nucleic acids include DNA, RNA, and fragments thereof. Nucleic acids may be derived from genomic

-13-

DNA, RNA, mitochondrial nucleic acid, chloroplast nucleic acid and other organelles with separate genetic material.

As used herein "nucleic acid" refers to polynucleotides such as deoxyribonucleic acid (DNA) and ribonucleic acid (RNA). The term should also be understood to include, as equivalents, derivatives, variants and analogs of either RNA or DNA made from nucleotide analogs, single (sense or antisense) and double-stranded polynucleotides. Deoxyribonucleotides include deoxyadenosine, deoxycytidine, deoxyguanosine and deoxythymidine. For RNA, the uracil base is uridine.

As used herein, "forced mass modulation" refers to methods provided herein that use deoxynucleotide analogs, modified nucleotide terminators, mass-labeled primers, mass-staggered primers and other such nucleotides, nucleic acids and analogs thereof, to unambiguously assign peak positions of mass fragments of oligonucleotides according to their base position, base identity, and target sequence from which the fragments arose. The method is used to sequence, detect or identify single oligonucleotide or plurality thereof. Hence the method is used, for example for multiplex sequencing and detection of nucleic acid molecules among mixtures thereof.

As used herein, "nucleotides" include, but are not limited to, the naturally occurring nucleoside mono-, di-, and triphosphates: deoxyadenosine mono-, di- and triphosphate; deoxyguanosine mono-, di- and triphosphate; deoxythymidine mono-, di- and triphosphate; and deoxycytidine mono-, di- and triphosphate (referred to herein as dA, dG, dT and dC or A, G, T and C, respectively). Nucleotides also include, but are not limited to, modified nucleotides and nucleotide analogs such as deazapurine nucleotides, *e.g.*, 7-deaza-deoxyguanosine (7-deaza-dG) and 7-deaza-deoxyadenosine (7-deaza-dA) mono-, di- and triphosphates, deuterio-deoxythymidine (deuterio-dT) mono-, di- and triphosphates, methylated nucleotides *e.g.*, 5-methyldeoxycytidine triphosphate, ¹³C/¹⁵N labelled nucleotides and deoxyinosine mono-, di- and triphosphate. For those skilled in the art, it will be clear that modified nucleotides and nucleotide analogs can be obtained using a variety of combinations of functionality and attachment positions.

-14-

As used herein, a complete set of chain-elongating nucleotides refers to the four different nucleotides or analogs thereof that hybridize to each of the four different bases comprising the nucleic acid template.

As used herein, the term "mass-matched nucleotides" refers to a set of
5 nucleotide analogs wherein each analog is of identical mass to each of the other analogs. For example, analogs of dA, dG, dC and dT can form a mass-matched nucleotide set, when each analog is selected to have the same molecular weight as the others in the set. Mass-matched nucleotide sets can be identified by selecting chemically modified derivatives of natural bases or by the use of a universal base
10 analog such as deoxyinosine or 5-nitroindole and 3-nitropyrrole (5-nitroindole and 3-intropyrrole can be in the dideoxy form) which can form base pairs with more than one of the natural bases. Others include, 3-methyl 7-propynyl iscarbostyrl, 5-methyl iscarbostyrl, and 3-methyl iscarbostyrl. As a result, oligonucleotides that contain such bases differ in molecular weight only as a function of length
15 thereof. Furthermore, incorporation of a single nucleotide(s) that is (are) not in the set renders such the oligonucleotide(s) readily identifiable by mass, particularly by spectrometric analysis.

As used herein, the term "pair-matched nucleotides" refers to a nucleotide set in which the nucleotide analogs are selected such that the total mass each base
20 pair is identical. For example, replacing dG with the nucleotide analog 7-deaza-dG forces the mass of each base pair, *i.e.*, (dA + dT) and (dC + 7-deaza-dG) to be identical. Exemplary pair-matched nucleotides, include, but are not limited to, 7-deaza-dA + phosphorothioate-dT ((312.2 + 320.2) = 632.4 Da) and 5-methyl-dC + dG ((303.2 + 329.2) = 632.4 Da); phosphorothioate-7-deaza-dA
25 + dU ((328.2 + 290.2) = 618.4 Da) and dC + dG = ((289.2 + 329.2) = 618.4 Da), and other such pairs that may be readily selected. Another exemplary set of mass-matched nucleotides with a molecular mass of 328.2: 7-deaza-dG, phosphorothioate-7-deaza-dA, 5-propynyl-dU and 5-cyanomethyl-2'-deoxycytidine.

As used herein, the term "nucleotide terminator" or "chain terminating
30 nucleotide" refers to a nucleotide analog that terminates nucleic acid polymer (chain) extension during procedures wherein a DNA template is being sequenced or replicated. The standard chain terminating nucleotides, *i.e.*, nucleotide terminators

-15-

include 2',3'-dideoxynucleotides (ddATP, ddGTP, ddCTP and ddTTP, also referred to herein as dideoxynucleotide terminators). As used herein, dideoxynucleotide terminators also include analogs of the standard dideoxynucleotide terminators, *e.g.*, 5-bromo-dideoxyuridine, 5-methyl-dideoxycytidine and dideoxyinosine are
5 analogs of ddTTP, ddCTP and ddGTP, respectively.

As used herein, "mass-matched terminators" refers to a set of nucleotide terminators that are selected such that each analog of ddA, ddG, ddC and ddT making up the mass-matched set has exactly the same molecular weight. Mass-matched terminator sets can be constructed by selecting chemically modified
10 derivatives of standard dideoxynucleotides or by the use of a universal dideoxynucleotide analog that form base pairs with more than one of the natural bases. Exemplary mass-matched nucleotides include, but are not limited to, 3-methyl 7-propynyl isocarbostyryl, 5-methyl iscarbostyryl and 3-methyl iscarbostyryl.

As used herein, the terms "oligonucleotide" or "nucleic acid" refer to single-
15 stranded and/or double-stranded polynucleotides such as deoxyribonucleic acid (DNA), and ribonucleic acid (RNA) as well as derivatives of either RNA or DNA into which nucleotide or dideoxynucleotide analogs have been incorporated. Also included in the term "nucleic acid" are analogs of nucleic acids such as peptide nucleic acid (PNA), phosphorothioate DNA, and other such analogs and derivatives.

As used herein, "nucleotide composition" or "base composition" refers to
20 the numerical ratio of the four nucleotide bases relative to each other in an oligonucleotide.

As used herein, a target nucleic acid refers to any nucleic acid of interest in a sample. It can contain one or more nucleotides. A target nucleotide sequence
25 refers to a particular sequence of nucleotides in a target nucleic acid molecule. Detection or identification of such sequence results in detection of the target and can indicate the presence or absence of a particular mutation or polymorphism.

As used herein, "partially duplex hairpin" refers to a partially self-complementary oligonucleotide, which forms intramolecular base-pairs within its
30 self-complementary region, leaving a "loop" of bases at one end of the molecule and a single-stranded "overhang" region at the other end. Thus, the oligonucleotide assumes a hairpin-like motif. "Blunt-ended hairpin structures", as referred to

-16-

herein, are similar to the partially duplex hairpin structures with the exception that they do not have a single-stranded "overhang" region.

As used herein, "base periodicity" or "period" (P_{base}) refers to the quasi-periodic distribution of the molecular weights of products obtained using Forced
5 Mass Modulation. The base periodicity results from either the mass of the mass-matched deoxynucleotide set, or the mass of the pair-matched deoxynucleotide set, or from the modified chain terminators depending on the embodiment implemented. The base sequence or nucleic acid molecule identity is encoded in the pattern (or detectable therein) in which the observed mass distribution deviates from absolute
10 regular periodicity.

As used herein, the "periodic reference mass" at base position "n" in any given oligonucleotide molecule, $M_{ref}[n]$, is defined as the sum of: (i) the mass of the primer (M_{primer}) used to sequence the DNA template using Forced Mass Modulation, (ii) the mass of the lightest nucleotide terminator (M_{light}), and, (iii) (n-1) multiple of
15 the base periodicity P_{base} .

As used herein, the "positional mass difference" or "mass shift" at base position "n" in any given oligonucleotide molecule, $M_{diff}[n]$, is defined as the distance in daltons between the observed peak, $M_{obs}[n]$, and the nth periodic reference mass.

20 As used herein, the "maximum mass shift" S_{max} is the maximum possible value of the positional mass difference, depending on the choice of mass-matched nucleotides and nucleotide terminators used in the implementation of Forced Mass Modulation. Accordingly, the maximum mass shift can be modulated by the choice of mass-matched nucleotides and nucleotide terminators.

25 As used herein, a "primer" refers to an oligonucleotide that is suitable for hybridizing, chain extension, amplification and sequencing. Similarly, a probe is a primer used for hybridization. The primer refers to a nucleic acid that is of low enough mass, typically about between about 5 and 200 nucleotides, generally about 70 nucleotides or less than 70, and of sufficient size to be conveniently used
30 in the methods of amplification and methods of detection and sequencing provided herein. These primers include, but are not limited to, primers for detection and sequencing of nucleic acids, which require a sufficient number nucleotides to form

-17-

a stable duplex, typically about 6-30 nucleotides, about 10-25 nucleotides and/or about 12-20 nucleotides. Thus, for purposes herein, a primer is a sequence of nucleotides contains of any suitable length, typically containing about 6-70 nucleotides, 12-70 nucleotides or greater than about 14 to an upper limit of about 5 70 nucleotides, depending upon sequence and application of the primer.

As used herein, the term "mass-labeled primers" refers to a set of primers that differ in mass by values that provide distinct and resolvable positional mass differences for each of the four termination reactions in an embodiment of Forced Mass Modulation. In this particular embodiment of Forced Mass Modulation, each 10 of the termination reactions for a given oligonucleotide is carried out separately using each of the mass-labeled primers, and the reaction products are combined prior to obtaining a mass spectrum.

As used herein, the term "mass-staggered primers" refers to the mass difference ("staggering" of the masses) between the primers used in multiplexed 15 sequencing using Forced Mass Modulation. For resolution of multiple sequences using this method, the differences between the masses of the primers should at least be equal to the maximum mass shift, and is generally greater than the maximum mass shift by at least 20 daltons to account for the finite width of each observed peak.

As used herein, reference to mass spectrometry encompasses any suitable 20 mass spectrometric format known to those of skill in the art. Such formats include, but are not limited to, Matrix-Assisted Laser Desorption/Ionization, Time-of-Flight (MALDI-TOF), Electrospray (ES), IR-MALDI (see, e.g., published International PCT application No.99/57318 and U.S. Patent No. 5,118,937), Ion Cyclotron Resonance 25 (ICR), Fourier Transform and combinations thereof. MALDI, particular UV and IR, are among the preferred formats.

As used herein, mass spectrum refers to the presentation of data obtained from analyzing a biopolymer or fragment thereof by mass spectrometry either graphically or encoded numerically.

As used herein, used herein, pattern with reference to a mass spectrum or 30 mass spectrometric analyses, refers to a characteristic distribution and number of signals (such peaks or digital representations thereof).

-18-

As used herein, signal in the context of a mass spectrum and analysis thereof refers to the output data, which the number or relative number of molecules having a particular mass. Signals include "peaks" and digital representations thereof.

- 5 As used herein, "mass spectrum division multiplexing" is an embodiment of Forced Mass Modulation in which unambiguous resolution of multiple sequences in a single spectrum is possible by judicious selection of mass staggered primers.

- As used herein, "analysis" refers to the determination of certain properties of a single oligonucleotide, or of mixtures of oligonucleotides. These properties
10 include, but are not limited to, the nucleotide composition and complete sequence of an oligonucleotide or of mixtures of oligonucleotides, the existence of single nucleotide polymorphisms between more than one oligonucleotide, the masses and the lengths of oligonucleotides and the presence of a molecule or sequence within molecule in a sample.

- 15 As used herein, "multiplexing" refers to the simultaneous determination of more than one oligonucleotide molecule, or the simultaneous analysis of more than one oligonucleotide, in a single mass spectrometric or other sequence measurement, *i.e.*, a single mass spectrum or other method of reading sequence.

- As used herein, "polymorphisms" refer to variants of a gene or an
20 oligonucleotide molecule that differ at more than one base position. In "single nucleotide polymorphisms", the variants differ at only a single base position.

- As used herein, amplifying refers to means for increasing the amount of a bipolymer, especially nucleic acids. Based on the 5' and 3' primers that are chosen, amplification also serves to restrict and define the region of the genome which is
25 subject to analysis. Amplification can be by any means known to those skilled in the art, including use of the polymerase chain reaction (PCR) etc. Amplification, *e.g.*, PCR must be done quantitatively when the frequency of polymorphism is required to be determined.

-19-

As used herein, "polymorphism" refers to the coexistence of more than one form of a gene or portion thereof. A portion of a gene of which there are at least two different forms, i.e., two different nucleotide sequences, is referred to as a "polymorphic region of a gene". A polymorphic region can be a single nucleotide, the identity of which differs in different alleles. A polymorphic region can also be several nucleotides in length. Thus, a polymorphism, e.g. genetic variation, refers to a variation in the sequence of a gene in the genome amongst a population, such as allelic variations and other variations that arise or are observed. Thus, a polymorphism refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. These differences can occur in coding and non-coding portions of the genome, and can be manifested or detected as differences in nucleic acid sequences, gene expression, including, for example transcription, processing, translation, transport, protein processing, trafficking, DNA synthesis, expressed proteins, other gene products or products of biochemical pathways or in post-translational modifications and any other differences manifested amongst members of a population. A single nucleotide polymorphism (SNP) refers to a polymorphism that arises as the result of a single base change, such as an insertion, deletion or change in a base.

A polymorphic marker or site is the locus at which divergence occurs. Such site may be as small as one base pair (an SNP). Polymorphic markers include, but are not limited to, restriction fragment length polymorphisms, variable number of tandem repeats (VNTR's), hypervariable regions, minisatellites, dinucleotide repeats, trinucleotide repeats, tetranucleotide repeats and other repeating patterns, simple sequence repeats and insertional elements, such as Alu. Polymorphic forms also are manifested as different mendelian alleles for a gene. Polymorphisms may be observed by differences in proteins, protein modifications, RNA expression modification, DNA and RNA methylation, regulatory factors that alter gene expression and DNA replication, and any other manifestation of alterations in genomic nucleic acid or organelle nucleic acids.

As used herein, "polymorphic gene" refers to a gene having at least one polymorphic region.

-20-

As used herein, "allele", which is used interchangeably herein with "allelic variant" refers to alternative forms of a gene or portions thereof. Alleles occupy the same locus or position on homologous chromosomes. When a subject has two identical alleles of a gene, the subject is said to be homozygous for the gene or
5 allele. When a subject has two different alleles of a gene, the subject is said to be heterozygous for the gene. Alleles of a specific gene can differ from each other in a single nucleotide, or several nucleotides, and can include substitutions, deletions, and insertions of nucleotides. An allele of a gene can also be a form of a gene containing a mutation.

10 As used herein, "predominant allele" refers to an allele that is represented in the greatest frequency for a given population. The allele or alleles that are present in lesser frequency are referred to as allelic variants.

As used herein, a subject, includes, but is not limited to, animals, plants, bacteria, viruses, parasites and any other organism or entity that has nucleic acid.

15 Among subjects are mammals, preferably, although not necessarily, humans. A patient refers to a subject afflicted with a disease or disorder.

As used herein, a phenotype refers to a set of parameters that includes any distinguishable trait of an organism. A phenotype can be physical traits and can be, in instances in which the subject is an animal, a mental trait, such as emotional
20 traits.

As used herein, "resolving power" of a mass spectrometer is the ion separation power of the instrument, *i.e.*, it is a measure of the ability of the mass spectrometer to separate peaks representing different masses. The resolving power R is defined as $m/\Delta m$, where m is the ion mass and Δm is the difference in mass
25 between two resolvable peaks in a mass spectrum.

As used herein, "assignment" refers to a determination that the position of a nucleic acid fragment indicates a particular molecular weight and a particular terminal nucleotide.

As used herein, "a" refers to one or more.

30 As used herein, "plurality" refers to two or more, up to an amount that is governed by the base periodicity, the maximum mass shift, and the resolving power of the mass spectrometer.

-21-

As used herein, an array refers to a pattern produced by three or more items, such as three or more loci on a solid support.

As used herein, "distinct" refers to a unique value of molecular weight, mass shift or period that is different from every other value of molecular weight,
5 mass shift or period in the measurement.

As used herein, "unambiguous" refers to the unique assignment of a particular oligonucleotide fragment according to the identity of its terminal base position and, in the event that a number of molecules are multiplexed, that the peak representing an oligonucleotide fragment can also be uniquely assigned to a
10 particular molecule.

As used herein, the symbols M_c , M_r , M_A and M_G are average molecular weights in daltons of the nucleotides deoxycytidine, thymidine, deoxyadenosine and deoxyguanosine, respectively, or of analogs thereof. M_{avg} , the average molecular weight of any given oligonucleotide is a function of the average molecular weights
15 of each of the nucleotides comprising the oligonucleotide, the numbers c , t , a and g of each nucleotide present in the oligonucleotide, the length of the oligonucleotide n' that is the sum of c , t , a and g , and the constant k that represents the mass of any other chemical groups on the molecule, such as terminal phosphates.

As used herein, N_{TOTAL} is the total number of possible base compositions for
20 an oligonucleotide of length n' .

As used herein, "peak separation" or "minimum peak separation" S refers to the minimum value of the distance between consecutive peaks in a mass spectrum that resolves a large number of oligonucleotides having the same lengths but different molecular weights, *i.e.*, different base compositions. The peak
25 separation, which can be tailored by careful selection of the nucleotide analogs incorporated into the oligonucleotide and by a mass spectrometer of desired resolving power, is usually a positive integer greater than one, and typically a positive integer greater than or equal to 3. For two oligonucleotides having the same length n' but different base compositions, their molecular weights will either
30 correspond to the same peak if the molecular weights are identical, or to two peaks separated at least by a value equal to the peak separation.

-22-

As used herein, L is the maximum number of allowed oligonucleotide masses for a given nucleotide set. It is directly proportional to the oligonucleotide length n' and the mass difference between the heaviest and lightest nucleotides in the set, and is inversely proportional to the peak separation.

5 As used herein, D refers to the average density of different base compositions per allowed mass value, given the set of all possible base compositions of an oligonucleotide of length n' .

As used herein, M_{heavy} refers to the mass of the heaviest nucleotide, nucleotide terminator or terminating base pair in daltons, depending on the specific
10 embodiment of Forced Mass Modulation being described.

As used herein, M_{light} refers to the mass of the lightest nucleotide, nucleotide terminator or terminating base pair in daltons, depending on the specific embodiment of Forced Mass Modulation being described.

As used herein, M_{primer} is the mass of the primer in daltons.

15 As used herein, $M_{\text{obs}}[n]$ is the observed mass of the sequencing reaction at the n th base position.

As used herein, $M_{\text{term}}[n]$ refers to the mass in daltons of the n th terminating nucleotide.

As used herein, L' is the theoretical upper limit on the number of sequences
20 that be multiplexed in a single mass spectrum. L' is directly proportional to the base periodicity P_{base} , and is inversely proportional to the maximum mass shift S_{max} .

As used herein, M_{duplex} is the mass in daltons of the fully duplex hairpin primer in the implementation of Forced Mass Modulation using pair-matched nucleotides.

25 As used herein, M_{ddm} is the mass in daltons of a dideoxy terminator that belongs to a set of mass-matched terminators.

As used herein, $M_{\text{seq}}[n]$ is the mass of the n th nucleotide past the priming site in the 3' to 5' direction in the target sequence, *i.e.*, the oligonucleotide whose sequence is being determined.

30 As used herein, "specifically hybridizes" refers to hybridization of a probe or primer only to a target sequence preferentially to a non-target sequence. Those of skill in the art are familiar with parameters that affect hybridization; such as

-23-

temperature, probe or primer length and composition, buffer composition and salt concentration and can readily adjust these parameters to achieve specific hybridization of a nucleic acid to a target sequence.

As used herein, a biological sample refers to a sample of material obtained
 5 from or derived from biological material, such as, but are not limited to, body fluids, such blood, urine, cerebral spinal fluid and synovial fluid, tissues and organs. Derived from means that sample can be processed, such as by purification or isolation and/or amplification of nucleic acid molecules.

As used herein, a composition refers to any mixture. It may be a solution,
 10 a suspension, liquid, powder, a paste, aqueous, non-aqueous or any combination thereof.

As used herein, a combination refers to any association between two or among more items.

As used herein, "kit" refers to a package that contains a combination and
 15 optionally instructions and/or reagents and apparatus for use with the combination.

Forced Mass Modulation for analysis of nucleic acid molecules

Time of flight analysis and drawbacks thereof

While time-of-flight mass spectrometry offers a number of advantages over conventional techniques such as gel electrophoresis, the peculiar relationship
 20 between the masses of the bases in DNA complicates the analysis of complex mixtures of oligonucleotides by mass spectrometry. For a given oligonucleotide, the average molecular weight, M_{avg} , is given by the following equation:

$$(i) \quad M_{avg} = k + cM_c + tM_t + aM_A + gM_G$$

where M_c , M_t , M_A , M_G are the average molecular weights of each of the four
 25 nucleotide bases (cytosine, thymine, adenine, guanine) and c , t , a , g represent the number of each base present in the oligonucleotide. The term k is a constant representing the mass of any other chemical groups on the molecule, such as terminal phosphates. Rearranging equation (i) to give the average molecular weight as a function of the length of the oligonucleotide in bases yields

$$30 \quad (ii) \quad M_{avg} = k + n'M_c + t(M_t - M_c) + a(M_A - M_c) + g(M_G - M_c)$$

where n' , the oligonucleotide length, is defined as

$$n' = c + t + a + g$$

-24-

Substituting the masses of the naturally occurring bases in DNA (to one-tenth dalton):

$$M_C = 289.2$$

$$M_T = 304.2$$

$$5 \quad M_A = 313.2$$

$$M_G = 329.2$$

into equation (ii) yields

$$(iii) \quad M_{avg} = k + 289.2n' + t(304.2 - 289.2) + a(313.2 - 289.2) + g(329.2 - 289.2),$$

10 which can be simplified to

$$(iv) \quad M_{avg} = k + 289.2n' + 15t + 24a + 40g$$

Close inspection of equation (iv) reveals that it is almost always possible to find two oligonucleotides of the same length but of different base composition whose average masses differ by only one dalton. For example, all 7-mers having a base composition of $A_2C_2G_2T$ have an average molecular weight of $(2167.4 + k)$, while all 7-mers with the base composition A_3CGT_2 have an average molecular weight of $(2166.4 + k)$. Since the following relation

$$(M_C + M_G) = (M_T + M_A) + 1$$

is always true for the naturally occurring bases in DNA, simply replacing one C and one G in an oligonucleotide with one A and one T will produce a new oligonucleotide exactly one dalton lighter. Many other "single-dalton difference" relations, such as

$$4M_A = (M_C + M_T + 2M_G) + 1$$

can readily be found for the naturally occurring bases.

25 Thus, the possibility always exists that two or more oligonucleotides of same length and different molecular weight (and, therefore, different base composition) will be too close in mass to be resolved by a time-of-flight instrument. Two oligonucleotides of same length but different molecular weight differ in base composition unless they are each composed of different nucleotide analogs, whereas two oligonucleotides of same length and same molecular weight can have either the same or different base compositions. This problem becomes increasingly severe with increasing oligonucleotide size, since the total number of possible base

-25-

compositions, N_{TOTAL} , scales as a cubic function of the oligonucleotide length n' , in bases:

$$(v) \quad N_{TOTAL} = \frac{(n' + 1)(n' + 2)(n' + 3)}{6}$$

5 The use of time-of-flight mass spectrometry in sequencing applications also poses several potential problems. The great drawback of sequencing by the Sanger method is that the molecular weights of the Sanger reaction products can appear virtually anywhere on the mass axis depending on the particular sequence being examined. As a result, the absolute mass of any single Sanger fragment has to
10 be measured with sufficient accuracy to calculate its distance from the masses from the fragments above and below it. Thus, determination of the identity of a single base depends on the accuracy of two separate mass measurements. Any error in a determination mass of a single fragment affects the accuracy of two bases in the sequence.

15 For longer sequences (30-50 bases), it may not be possible to determine the mass difference between adjacent peaks with sufficient accuracy to unambiguously determine base identity. This is particularly a problem for the nucleotides A and T, which differ in mass only by nine daltons. The problem is addressed by resolving each of the four termination reactions in a separate mass spectrum. In this case
20 each peak functions essentially as binary signal indicating the presence of a base at a particular position, much as in conventional electrophoretic sequencing. Using separate spectra, however, increases read accuracy but at the expense of increasing the number of required mass measurements by a factor of four.

25 It is possible to resolve two target sequences by the Sanger method in a single mass spectrum, provided that all products of the sequencing reactions have unique and resolvable masses, and multiplex methods using mass modified bases have been developed. But, where two or more reaction products have the same mass, then unambiguous reconstruction of the two target sequences is not possible (see, *e.g.*, Figures 1c-e). In addition, there is no way to determine *a priori* which
30 observed masses belong to a particular sequence. In practice, this means that multiplexed Sanger sequencing by mass spectrometry can be difficult. The

-26-

methods provided herein resolve these problem and provide a way to determine *a priori* which masses are associated with extension of a particular primer.

Forced Mass Modulation

As noted above, Forced Mass Modulation refers to methods provided herein
5 that permit unambiguously assign peak positions (or masses) to mass fragments of
oligonucleotides according to their base position, base identity, and target sequence
from which the fragments arose. The methods use deoxynucleotide analogs,
modified nucleotide terminators, mass-labeled primers, mass-staggered primers and
other such nucleotides, nucleic acids and analogs thereof to provide a means for
10 deconvoluting complex mass spectra or output from other mass determining
techniques. These methods permit deconvolution of highly multiplexed nucleic acid
reaction mixtures for sequencing methods and detection methods that include a
step of primer extension. In practicing these methods, primers are extended using
mass-matched nucleotides and chain terminators (or in some embodiments mass
15 where it is only necessary to detect incorporation (or the absence of incorporated)
mass-matched terminators and optionally mass-matched chain extending
nucleotides). Because the sequence and/or molecular mass of a primer is known,
and the extended nucleotides have the same molecular mass, a periodicity in
molecular mass that is a function of molecular weight of the selected mass matched
20 nucleotide(s) results.

As described in more detail below, for sequencing reactions using chain
terminators, the deviation from the periodicity results from incorporation of a chain
terminator. The deviation is a function of the particular terminator incorporated.
For detection methods, incorporation of a terminator will indicate the presence of
25 a mutation (if the terminator is selected to pair with the first mutated nucleotide.
Any shift from periodicity will indicate the presence of the mutation. These
methods, thus provide a simple, reliable way to detect the presence of a mutations
or target nucleotide(s) in a sequence and to sequence nucleic acids. Forced mass
modulation can be used with any method, such as mass spectrometry and gel
30 electrophoresis, that relies on molecular weight as an output. Mass spectrometry
is exemplified herein.

-27-

The methods, designated Forced Mass Modulation methods, provided herein, are implemented by suitable selection of nucleotides and/or chain terminators, such as by the use of deoxynucleotide analogs, modified nucleotide terminators and mass-labeled primers in one or more reactions. Forced Mass Modulation can be used to simultaneously sequence or detect large numbers, such twenty-five or more) oligonucleotides, with a high degree of resolution and accuracy. It can also be used to simplify the analysis of closely related sequence variants, as is required in the detection and scoring of nucleotide polymorphisms, including single nucleotide polymorphism (SNPs) and for other genotypical analyses. Forced Mass Modulation greatly improves the use of mass spectrometry for nucleic acid analyses. Nearly every application relies on mass measurements that can benefit in increased accuracy and in a reduction of the number of required spectra. Another advantage of Forced Mass Modulation is the number of different ways in which it can be implemented, allowing it to be tailored to particular experimental or Instrumental limitations.

For example, compared to the conventional Sanger methods, Forced Mass Modulation, provides increased accuracy, simplified interpretation of mass data, and the ability to use a single mass spectrum for the unambiguous resolution of several distinct nucleic acid molecules. For mass spectrometry applications, the methods provide unambiguous assignment of peak positions of mass fragments of oligonucleotides according to their base position, base identity, and target sequence from which the fragments arose. Thus, the methods herein are advantageously used for multiplexing, in which a plurality of reactions are run in a single reaction (single pot). Forced Mass Modulation, exemplified with reference to sequencing methods, such as PROBE, can also be adapted to detection methods in which a primer is extended.

In Forced Mass Modulation in which a primer is extended with mass-matched nucleotides, for examples, the molecular weights of extended nucleic acid chains, such as sequencing reaction products, are constrained since all extension products from the same primer will have a molecular weight that differs either by the length of the extension and the chain terminator. As a result, the extension products assume a quasi-periodic distribution on the mass axis with a

-28-

predetermined *base periodicity*. For sequencing, the base sequence itself is encoded in the pattern in which the observed mass distribution deviates from absolute regular periodicity. Since the base periodicity will always be known *a priori*, since the primer is known, each peak in the observed mass spectrum can be matched unambiguously to a unique nucleotide position in the target sequence. The initiating primers fix each set of nested fragments or extended products, and the use of mass-matched nucleotides creates the periodicity.

As demonstrated by the Examples below, the method is advantageous for numerous applications including sequencing and a variety of detection methods, including primer oligo base extension (PROBE) (see, *e.g.*, U.S. application Serial No. 6,043,031; allowed U.S. application Serial No. 09/287,679; and 6,235,478) that use mass spectrometry to distinguish between extended primers. If the base compositions of the target oligonucleotides are known *a priori* then it is possible to select a nucleotide set that produces oligonucleotide masses that are distinct and resolvable for any particular instrument or application.

Conversely, it is also possible to select a nucleotide set that restricts specific oligonucleotides to have the same mass, regardless of a change in base composition. The strategy of restricting specific oligonucleotides to have the same mass can be used to separate more than one oligonucleotide population of different lengths by restricting all oligonucleotides of a particular length to the same molecular weight, irrespective of differences in base composition.

The oligonucleotide analysis or sequencing in methods provided herein can be accomplished by one of several methods employed in the art for the synthesis, resolution and/or detection of nucleic acids. Depending on the embodiment implemented, modified nucleotides can be incorporated into the oligonucleotides by chemical (*Oligonucleotides and Analogues: A Practical Approach*, F. Eckstein, ed., IRL Press Oxford, 1991) or enzymatic (F. Sanger et al., *Proc. Natl. Acad. Sci. USA* 74:5463-67, 1977) synthesis. Extension products or truncated products of the oligonucleotides to be sequenced can be obtained using chemical (A.M. Maxam and W. Gilbert, *Proc. Natl. Acad. Sci. USA* 74:560-64, 1977) or enzymatic (F. Sanger et al., *Proc. Natl. Acad. Sci. USA* 74:5463-67, 1977) methods.

-29-

For the resolution and detection of target nucleic acids any mass determination method, such as, but are not limited to, chromatography, gel electrophoresis, capillary zone electrophoresis and mass spectrometry, is used. Mass spectrometric formats, include, but are not limited to, are matrix assisted laser desorption ionization (MALDI), electrospray (ES), ion cyclotron resonance (ICR) and Fourier Transform. For ES, the samples, dissolved in water or in a volatile buffer, are injected either continuously or discontinuously into an atmospheric pressure ionization interface (API) and then mass analyzed by a quadrupole. The generation of multiple ion peaks which can be obtained using ES mass spectrometry can increase the accuracy of the mass determination. Even more detailed information on the specific structure can be obtained using an MS/MS quadrupole configuration

In MALDI mass spectrometry, various mass analyzers can be used, *e.g.*, magnetic sector/magnetic deflection instruments in single or triple quadrupole mode (MS/MS), Fourier transform and time-of-flight (TOF) configurations as is known in the art of mass spectrometry. For the desorption/ionization process, numerous matrix/laser combinations can be used. Ion-trap and reflectron configurations can also be employed.

Pair-matched nucleotide-based methods

Forced Mass Modulation can be implemented using a deoxynucleotide set in which the mass of each *base pair* is identical, termed a *pair-matched* nucleotide set. A pair-matched nucleotide set can easily be formed, for example, by replacing dG (329.2 Da) in the set of naturally occurring nucleotides with 7-deaza-dG (328.2 Da). This forces the mass of each base pair to be 617.4 daltons:

$$\begin{aligned} (dA + dT) &= (313.2 + 304.2) = 617.4 \text{ Da} \\ (dC + 7\text{-deaza-dG}) &= (289.2 + 328.2) = 617.4 \text{ Da} \end{aligned}$$

Many other pair-matched sets are possible using available nucleotide analogs. For this embodiment, the target DNA sequence can be composed *entirely* of the pair-matched nucleotide set. This can be accomplished by amplifying the target DNA sequence by PCR using the pair-matched nucleotide set prior to the sequencing reaction.

A further requirement for this embodiment of Forced Mass Modulation is that the mass of each *terminating* base pair is unique and resolvable. The

-30-

standard dideoxy terminators therefore cannot be used with the pair-matched nucleotide set described above, because the masses of all terminating base pairs are identical at 601.4 daltons, except ddG:dC, which is 602.4 daltons. For the sake of clarity in this example, it is assumed that a set of mass-matched

5 terminators is available ("ddM," defined as set of chain-terminating nucleotides that have exactly the same molecular weight $ddA = ddC = ddG = ddT$). If the mass of ddM is arbitrarily chosen to be 500 daltons, then the masses of the terminating base pairs are as follows:

	<u>Terminating Base Pair</u>	<u>Mass (Da)</u>
10	<i>ddM: dC</i>	<i>789.2</i>
	<i>ddM: dT</i>	<i>804.2</i>
	<i>ddM: dA</i>	<i>813.2</i>
	<i>ddM: 7-deaza-dG</i>	<i>828.2</i>

In practice it is also possible to implement Forced Mass Modulation using a set

15 of terminators that have different masses, this is discussed in detail below.

Exemplary embodiments in which the mass shift is obtained using pair-matched nucleotides, where the mass of each nucleotide base-pair is selected so that the masses of all pairs are identical, are described in Example 4. In one embodiment thereof, the following steps are performed: (i) the target nucleic

20 acid is copied or amplified by a method such as PCR in the presence of the pair-matched nucleotide set prior to the sequencing or detection reaction; (ii) the target nucleic acid is denatured, and a partially duplex hairpin primer is annealed and ligated to the single-stranded template; (iii) the primer is extended in the presence of chain terminating nucleotides and pair-matched nucleotides to

25 produce extension products; (iv) the masses of the extension products follow a periodic distribution that is determined by the mass of the pair-matched nucleotide set, and, (v) the target nucleic acid is detected by virtue of its molecular weight or its sequence is determined from the mass shift of each extension product from its corresponding periodic reference mass.

-31-

In embodiments described above, the extending bases are pair matched. the mass of each terminating base pair is unique and resolvable, so that the mass shifts corresponding to each terminating base pair are unique. The nucleotide terminators are optionally mass-matched or can be of distinct masses as long as distinct values of mass shift are obtained for each terminating base pair.

In another embodiment, the extension products are treated to produce blunt-ended double-stranded extension products by methods known to those of skill in the art, such as the use of single-strand specific nucleases. In an aspect of this embodiment, a plurality of target nucleic acids can be multiplexed in a single reaction by annealing each target nucleic acid to a primer of distinct molecular weight. The primers can be selected to differ in molecular weight by a value that is greater than the maximum mass shift, *i.e.*, the difference in molecular weight between the heaviest and the lightest nucleotide terminating base pairs. Since double stranded nucleic acid can be analyzed, the effective sequence read is halved relative to the embodiment employing mass-matched nucleotides, but the number of molecules that can be multiplexed is doubled, due to the increase in period (the value of the mass of a base pair, rather than a single mass-matched nucleotide). In exemplary embodiments, about 14 to about 50 sequences are multiplexed. In detection embodiments, about 50 or more molecules can be simultaneously multiplexed since only a single terminating base pair is added in the extension reaction.

In another embodiment, the chain termination reactions can each be carried out separately using a standard nucleotide terminator, pair-matched nucleotides, and mass-labeled primers, if modified nucleotide terminators which are either mass-matched or provide distinct values of mass shift for each terminating base pair are not available. The reactions can be pooled prior to detection or sequence analysis. In one embodiment, the mass-labeled primers can have distinct values of molecular weight that give rise to unique values of mass shift or positional mass difference for each terminating base.

-32-

Optimizing the mass spectrometric analysis of oligonucleotide mixtures

In another method provided herein, nucleotide analogs are used to restrict the possible values of molecular weights that an oligonucleotide can possess relative to other oligonucleotides of the same length. The nucleotide analogs can be incorporated into the oligonucleotides using any suitable method, such as automated DNA synthesis (*Oligonucleotides and Analogues: A Practical Approach*, F. Eckstein, ed., IRL Press Oxford, 1991) or by enzymatic replication using a polymerase and the requisite nucleotides and nucleotide analogs.

For example, any two oligonucleotides with the same length n' with different base compositions can either 1) have exactly the same average molecular weight, or 2) have molecular weights no closer than a minimum value called the *peak separation*. In most cases, the peak separation will be a positive integer greater than one, but fractional values are theoretically possible.

To illustrate an exemplary implementation of this method, the average molecular weight of the nucleotide analog 7-deaza-dG (328.2 daltons) can be substituted for M_G , into equation (ii) above, which defines M_{avg} as a function of the length " n " of the oligonucleotides in bases, as follows:

(ii) $M_{avg} = k + n'M_c + t(M_T - M_c) + a(M_A - M_c) + g(M_G - M_c)$, where M_c , M_T , M_A , M_G are the average molecular weights of each of the four nucleotide bases (cytosine, thymine, adenine, guanine); c , t , a , g represent the number of each base present in the oligonucleotide, the sum thereof, *i.e.*, $c + t + a + g = n'$, the total oligonucleotide length in bases; and the term k is a constant representing the mass of any other chemical groups on the molecule, such as terminal phosphates.

Substituting the masses of the naturally occurring bases dC, dT and dA in DNA (to one-tenth dalton), and of 7-deaza-dG,

$$M_c = 289.2$$

$$M_T = 304.2$$

$$M_A = 313.2$$

$$M_G = 328.2$$

and following simplification, the equation reduces to:

$$M_{avg} = k + 289.2n' + 15t + 24a + 39g$$

-33-

Extracting the common factor from the last three terms yields

$$(vi) M_{avg} = k + 289.2n' + (5t + 8a + 13g) \times 3$$

In this example, the minimum peak separation is three daltons. It is not possible to identify or detect two oligonucleotides of the same length with different molecular weights that are closer than three daltons. Oligonucleotides with average masses closer than three daltons the oligonucleotides are detected if they are of different lengths.

As a second example, Mr can be substituted with the molecular weight of a hypothetical nucleotide analog whose mass is 305.2 into equation (ii), yielding

$$10 \quad M_{avg} = k + 289.2n' + 16t + 24a + 40g$$

Extracting the common integer factor from the last three terms yields

$$(vii) M_{avg} = k + 289.2n' + (2t + 3a + 5g) \times 8$$

for a minimum peak separation of eight daltons. Thus, appropriate selection of nucleotide analogs permits construction of nucleotide sets that provides sufficient peak separation for adequate resolution by mass, such as in a time-of-flight mass spectrometer. The trade-off for a greater peak separation is a greater number of base compositions that have exactly the same mass for a given oligonucleotide length. The maximum number of allowed oligonucleotide masses, L, for a given nucleotide set, is given by

$$20 \quad (viii) L = \frac{n'(M_{heavy} - M_{light})}{S} + 1,$$

where n' is the oligonucleotide length in bases, S is the peak separation, M_{light} the mass of the lightest nucleotide in the set, M_{heavy} is the mass of the heaviest nucleotide in the set. The number of allowed oligonucleotide masses scales in direct proportion to the base length and inversely with the peak separation, but not all possible mass values will be represented for a given oligonucleotide length, particularly for small n. The average density of different base compositions per allowed mass value, D, can be obtained by dividing equation (v) by (viii)

$$30 \quad D = \frac{N_{TOTAL}}{L}$$

which expands into

$$(ix) \quad D = \frac{S(n' + 1)(n' + 2(n' + 3))}{6(n'(M_a - M_c) + S)}$$

-34-

using a typical nucleotide set with G as the heaviest base and C as the lightest. The density function scales in direct proportion to the peak separation and as a quadratic function of the oligonucleotide length in bases. In practice, the average density of base compositions per *allowed* mass value predicated by
5 equation (ix) will be somewhat lower than the actual density of base compositions per *observed* mass value, because not all allowed masses will always be represented. The Examples describe implementation of the methods for sequencing.

System and Software method for Force Mass Modulation

10 Also provided are systems that automate the methods for determining a nucleotide sequence of a target nucleic acid or the detection methods provided herein using a computer programmed for identifying the sequence or target nucleic acid identity based upon the methods provided herein. The methods
15 herein can be implemented, for example, by use of the following computer systems and using the following calculations, systems and methods.

An exemplary automated testing system contains a nucleic acid workstation that includes an analytical instrument, such as a gel electrophoresis apparatus or a mass spectrometer or other instrument for determining the mass of a nucleic acid molecule in a sample, and a computer capable of
20 communicating with the analytical instrument (see, *e.g.*, copending U.S. application Serial Nos. 09/285,481, 09/663,968 and 09/836,629; see, also International PCT application No. WO 00/60361 for exemplary automated systems). In an exemplary embodiment the computer is an IBM compatible computer system that communicates with the instrument using a known
25 communication standard such as a parallel or serial interface.

-35-

For example, systems for analysis of nucleic acid samples are provided. The systems include a processing stations that performs a forced mass modulation chain extension reaction; a robotic system that transports the resulting products from the processing station to a mass measuring station, 5 where the masses of the products of the reaction are determined; and a data analysis system, such as a computer programmed to identify nucleotides using forced mass modulation data, that processes the data from the mass measuring station to identify a nucleotide or plurality thereof in a sample or plurality thereof. The system can also include a control system that determines when 10 processing at each station is complete and, in response, moves the sample to the next test station, and continuously processes samples one after another until the control system receives a stop instruction.

The computer can be part of the instrument or another system component or it can be at a remote location. A computer system located at a 15 site distant from the instrument can communicate with the instrument, for example, through a wide area network or local area communication network or other suitable communication network. The system with the computer is programmed to automatically carry out steps of the methods herein and the requisite calculations. For embodiments that use mass-matched 20 deoxyribonucleotides, a user enters the primer sequence or primer mass, the periodic reference mass and mass of an individual mass-matched deoxynucleotide. These data can be directly entered by the user from a keyboard or from other computers or computer systems linked by network connection, or on removable storage medium such as a CD-ROM, minidisk (MD), 25 DVD, floppy disk or other suitable storage medium. Next the user causes execution software that operates the system in which the mass spectrum of the extension products is generated. The Forced Mass Modulation software performs the steps of obtaining the masses of the fragments generated by the sequencing reaction and measured by the analytical instrument, and determining the identity 30 of a nucleotide at any base position or the positional mass difference. The identity of the nucleotide at each base position is determined by comparing the

-36-

calculated $M_{diff}[n]$ values to a database of previously calculated values of M_{diff} for each of the chain terminating nucleotides.

$$M_{diff}[n] = M_{obs}[n] - M_{PR}[n],$$

where:

$$(i) \quad M_{PR}[n] = (M_{primer} + M_{light}) + (n - 1) P_{base},$$

in which n is the base position, $M_{PR}[n]$ is the n^{th} periodic reference mass, M_{primer} is the mass of the primer, M_{light} is the mass of the lightest nucleotide terminator and P_{base} is the base periodicity in daltons. The observed masses of the sequencing reaction products are given by the following equation:

$$(ii) \quad M_{obs}[n] = M_{primer} + (n - 1) P_{base} + M_{term}[n],$$

where n is the base position, $M_{obs}[n]$ is the n^{th} observed mass, P_{base} is the base periodicity, and $M_{term}[n]$ is the mass of the n^{th} terminating nucleotide in daltons. The positional mass differences for the sequence can be obtained by subtracting equation (i) from equation (ii) and evaluating at every base position n :

where $M_{diff}[n]$ is the n^{th} positional mass difference. This relation simplifies to:

$$(iii) \quad M_{diff}[n] = M_{term}[n] - M_{light}.$$

Hence, the periodicity is determined by the mass of the mass-matched nucleotide and the shift is the difference in location of a peak resulting from the chain terminator. For example, in Figure 2, the lightest terminator is ddC, and the differential is 0 for C, 40 for G, 34 for A, 15 for T. The selected mass matched nucleotide has a mass of 310 Da. The primer in Figure 2a has a mass of 3327 Da and the first peak would be at 3600 if the first nucleotide in the extension product were C (0 shift). Since the first peak is at 3640, the shift is 40 Da. Therefore the first nucleotide is G, corresponding to a shift from the periodicity of 310 Da generated by the mass-matched nucleotides.

Detection methods

The methods herein may be used with any method for detection of nucleic acids based on molecular mass known to those of skill in the art, particularly methods in which a primer is extended. Such methods are modified by extending using mass matched nucleotides and/or chain terminators in extension reactions. Alternatively, or additionally, amplification reactions may be performed using mass-matched nucleotides or pair-matched sets of nucleotides.

-37-

These methods can be readily multiplexed using the methods and nucleic acid molecules provided herein.

Detection methods and protocols, including those that rely on mass spectrometry (see, *e.g.*, U.S. Patent No. 6,194,144; 6,225,450; 5,691,141; 5,547,835; 6,238,871; 5,605,798; 6,043,031; 6,197,498; 6,235,478; 6,221,601; 6,221,605; International PCT application No. WO 99/31273, International PCT application No. WO 98/20019), can be modified for use with the methods herein by using mass-matched nucleotides for extension or pair matched duplexes for hybridization reactions.

Among the methods of analysis herein are those involving the primer oligo base extension (PROBE) reaction with mass spectrometry for detection. In such reactions, the primer will be extended by mass-matched nucleotides. The methods herein are designed for multiplexing so that a plurality of different primers can be extended at different loci in the same reaction. The PROBE method uses a single detection primer followed by an oligonucleotide extension step to give products, which can be readily resolved by mass spectrometry, and, in particular, MALDI-TOF mass spectrometry. The products differ in length depending on the presence or absence of a polymorphism. In this method, a detection primer anneals adjacent to the site of a variable nucleotide or sequence of nucleotides and the primer is extended using a DNA polymerase in the presence of one or more dideoxy NTPs and, optionally, one or more deoxy NTPs. The resulting products are resolved by MALDI-TOF mass spectrometry. The mass of the products as measured by MALDI-TOF mass spectrometry makes possible the determination of the nucleotide(s) present at the variable site. Use of primers containing mass-matched bases increases the resolving power of the reaction and permit simultaneous detection of a plurality of mutations (polymorphisms).

These methods can be automated (see, *e.g.*, copending U.S. application Serial No. 09/285,481 and published International PCT application No. PCT/US00/08111, which describes an automated process line) and performed in a system that includes a computer programmed for analysis of the mass data as described above.

-38-

The analyses can be performed on chip based formats in which the target nucleic acids or primers are linked to a solid support, such as a silicon or silicon-coated substrate, preferably in the form of an array. Generally, when analyses are performed using mass spectrometry, particularly MALDI, small nanoliter
5 volumes of sample are loaded on, such that the resulting spot is about, or smaller than, the size of the laser spot. It has been found that when this is achieved, the results from the mass spectrometric analysis are quantitative. The area under the signals in the resulting mass spectra are proportional to concentration (when normalized and corrected for background). Methods for
10 preparing and using such chips are described in U.S. Patent No. 6,024,925, co-pending U.S. application Serial Nos. 08/786,988, 09/364,774, 09/371,150 and 09/297,575; see, also U.S. application Serial No. PCT/US97/20195, which published as WO 98/20020. Chips and kits for performing these analyses are commercially available from SEQUENOM under the trademark MassARRAY.
15 MassArray relies on the fidelity of the enzymatic primer extension reactions combined with the miniaturized array and MALDI-TOF (Matrix-Assisted Laser Desorption Ionization-Time of Flight) mass spectrometry to deliver results rapidly. It accurately distinguishes single base changes in the size of DNA fragments associated with genetic variants without tags.

20

The following Examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

EXAMPLE 1

Forced Mass Modulation using Mass-Matched Deoxynucleotides

25 For this implementation, a set of nucleotide analogs for the four bases in DNA are selected (Amersham Pharmacia Biotech) such that each base has exactly the same molecular weight, termed a *mass-matched* deoxynucleotide set. This is achieved by judiciously choosing chemical modifiers of the existing bases or by the using a universal base analog such as deoxyinosine, which can
30 form base pairs with more than one of the natural bases. For this example, the mass of each deoxynucleotide ("dN") in the mass-matched set has the arbitrarily selected value of 310 daltons, but any other value suffices. The sequencing

-39-

reaction is performed as follows: 1) a primer is annealed to the target to be sequenced; 2) the resulting structure is subjected to a extension reaction using a suitable polymerase in the presence of the mass-matched nucleotide set and the four standard dideoxynucleotide terminators. The products and molecular masses of such a reaction are shown with a simulated mass spectrum in Figure 2a. The base periodicity is the mass of dN, or 310 daltons. The identity of a nucleotide at any base position is given by the *positional mass difference*, defined as the distance in daltons between the observed peak and the nearest *periodic reference mass*, which occurs every 310 daltons. In this example, the first periodic reference mass is defined as the (primer mass + ddC), or $(3327 + 273) = 3600$ daltons. The second periodic reference mass would be 3600 plus the base periodicity or $(3600 + 310) = 3910$, and so on. Expressed in terms of the base position n :

$$(i) \quad M_{PR}[n] = (M_{primer} + M_{light}) + (n - 1) P_{base},$$

where n is the base position, $M_{PR}[n]$ is the n^{th} periodic reference mass, M_{primer} is the mass of the primer, M_{light} is the mass of the lightest nucleotide terminator and P_{base} is the base periodicity in daltons. The observed masses of the sequencing reaction products are given by the following equation:

$$(ii) \quad M_{obs}[n] = M_{primer} + (n - 1) P_{base} + M_{term}[n],$$

where n is the base position, $M_{obs}[n]$ is the n^{th} observed mass, P_{base} is the base periodicity, and $M_{term}[n]$ is the mass of the n^{th} terminating nucleotide in daltons. The positional mass differences for the sequence can be obtained by subtracting equation (i) from equation (ii) and evaluating at every base position n :

$$M_{diff}[n] = M_{obs}[n] - M_{PR}[n],$$

where $M_{diff}[n]$ is the n^{th} positional mass difference. This relation simplifies to:

$$(iii) \quad M_{diff}[n] = M_{term}[n] - M_{light}.$$

Inspection of equation (iii) reveals that M_{diff} can only take on four distinct values, each corresponding to a different nucleotide terminator:

$$M_{diff}["ddC"] = (273.2 - 273.2) = 0$$

$$M_{diff}["ddT"] = (288.2 - 273.2) = 15$$

$$M_{diff}["ddA"] = (297.2 - 273.2) = 24$$

$$M_{diff}["ddG"] = (313.2 - 273.2) = 40.$$

-40-

Hence, the identity of the nucleotide at every base position in the target sequence can be determined by comparing each calculated positional mass difference with the values in the table above. Since the values that M_{diff} can assume depend only on the choice of nucleotide terminators used in the sequencing reaction, it is possible to tailor the positional mass differences so that they are resolvable for any particular mass spectrometer. For example, replacing the terminator ddT with its analog 5-bromo-dideoxyuridine (353.1 daltons) yields a positional mass difference of $(353.1 - 273.2) = 79.9$ Da for termination at T positions in the target sequence. This type of nucleotide substitution can be particularly valuable for lower-resolution mass spectrometers, as it is possible to maintain the sequence read accuracy without requiring any additional mass spectra.

Further inspection of equation (iii) reveals that each observed mass value can be at most 40 daltons heavier than the nearest periodic reference mass. This limit is termed the *maximum mass shift* and is defined as the mass difference between the heaviest nucleotide terminator and the lightest. Resolving a second target sequence by Forced Mass Modulation with the standard dideoxy terminators is possible in a single spectrum so long as the primer for the second sequence is at least 40 daltons heavier (the maximum mass shift) than the primer for the first sequence, thus insuring that the peaks for each sequence never overlap in mass.

In practice, it is recommended most mass spectrometric formats that the second primer is at least about 60 daltons heavier than the first primer, as each observed peak will have a finite width. Figure 2b shows a target second sequence resolved on the same mass spectrum shown in Figure 2a, using a primer heavier by 77 daltons. The peaks corresponding to the reaction products from the first target sequence can fall within the shaded regions of the spectrum in Figure 2b, which can never intersect peaks from the second target sequence. Unambiguous resolution of both sequences is possible in this arrangement because each peak can be uniquely assigned to a nucleotide, a base position, and a target sequence. This method is designated *Mass Spectrum Division Multiplexing* herein, and it is implemented using *mass-staggered primers*. Figure

-41-

3 shows four different sequences resolved in a single spectrum using a set of mass-staggered primers that are separated in mass by integer multiples of 77 daltons (77, 154, and 231 daltons).

The theoretical upper limit on the number of sequences that can be multiplexed in a single mass spectrum is given by the following equation:

$$(iv) L' = \frac{P_{base}}{S_{max}},$$

where L' is the upper limit, P_{base} is the base periodicity, and S_{max} is the maximum mass shift in daltons. For the nucleotide set and terminators used in this example, $L = (310 / 40) \approx 7.75$, or approximately seven. Increasing the number of sequences that can be multiplexed in a single spectrum, can be achieved by implementing one or both of an increase in the base periodicity, and a reduction of the maximum mass shift. The base periodicity can be increased by choosing a mass-matched nucleotide set that has a higher molecular weight for dN. It is simpler to lower the maximum mass shift by careful use of the nucleotide terminators and their analogs. For example, if the sequencing reactions were performed using only the terminators ddC, ddT, and ddA, then the maximum mass shift becomes (mass of ddA - mass of ddC) = (297 - 273) = 24 Da. In this case the upper limit on the number of sequences that can be multiplexed is $L = (310 / 24) \approx 12.92$, or approximately twelve. In situations where complete sequence information is not required, such as diagnostic sequencing, a great reduction in the number of required spectra can be realized by using fewer than four nucleotide terminators. If the sequencing reaction is performed using only a *single* nucleotide terminator, the maximum mass shift becomes identically zero, and the number of sequences that can be multiplexed in a single spectrum is limited only by the absolute resolution of the mass spectrometer in question. If a given mass spectrometer has an absolute resolution of 12 Da in the mass range of the sequencing reaction products, then the maximum number of sequences that can be multiplexed is given by $L = (310 / 12) \approx 25.83$, or approximately twenty-five.

-42-

EXAMPLE 2**Forced Mass Modulation using Pair-Matched Deoxynucleotides**

Implementation of Forced Mass Modulation using pair-matched nucleotides is shown in Figure 4. The basic requirement for this method is that the sequencing reaction products can be analyzed as double-stranded structures. Briefly, the steps in the reaction are as follows: 1) A partially duplex hairpin primer with a 3' overhang and a 5' phosphate group is annealed and ligated to the single stranded target sequence. 2) The resulting partially duplex structure is subjected to a sequencing reaction using the pair-matched nucleotide set described above along with the set of mass-matched terminators (ddM). 3) The products from the sequencing reaction are exposed to a strict single strand-specific nuclease that results in the production of blunt-ended hairpin structures ready for analysis by mass spectrometry. Figure 5 shows the products and molecular masses of the nuclease digestion along with a simulated mass spectrum.

Because the reaction products are double-stranded, they are forced to assume a quasi-periodic distribution with a base periodicity of 617.4 daltons. The shaded regions on the spectrum shown in Figure 5 indicate the allowed mass ranges that can be occupied by the reaction products. The first periodic reference mass is at 10360 Da, which is the mass of the fully duplex hairpin primer plus a ddM:dC base pair. Expressing the periodic reference masses in terms of the base position n yields:

$$(x) \quad M_{PR}[n] = (M_{duplex} + M_{light} + M_{ddM}) + (n - 1) \times P_{base}$$

Where $M_{PR}[n]$ is the n^{th} periodic reference mass, M_{duplex} is the mass of the fully duplex primer, M_{light} is the mass of the lightest deoxynucleotide in the target, P_{base} is the base periodicity, and M_{ddM} is the mass of ddM in daltons. The observed masses of the sequencing reaction products are given by the following equation:

$$(xi) \quad M_{obs}[n] = M_{duplex} + M_{ddM} + (n - 1) \times P_{base} + M_{targ}[n],$$

where n is the base position, $M_{obs}[n]$ is the n^{th} observed mass, and $M_{targ}[n]$ is the mass of the n^{th} nucleotide in the target sequence past the priming site in the 3' -> 5' direction.

-43-

In contrast to the mass-matched nucleotide set implementation that provides the sequence complementary to the template strand read in the 5' -> 3' direction, the pair-matched nucleotide set implementation described herein directly reads the template strand in the 3' -> 5' direction. The positional mass differences for this implementation are the same as those in Example 1, except that the mass difference corresponding to a termination on dG is 39 as opposed to 40 daltons, because 7-deaza-dG is exactly one dalton lighter than dG. Since double stranded DNA can be analyzed for this method to work, the effective sequence read length is halved, although the number of sequences that can be multiplexed is doubled, due to the increase in the base periodicity.

As a demonstration of Forced Mass Modulation implemented without using mass-matched terminators, the positional mass differences for the above example using the following set of nucleotide terminators is calculated as follows:

<u>Terminator</u>	<u>Nucleotide Analog</u>	<u>Mass</u>	<u>Base Pairing</u>	<u>Mass of Base Pair</u>
T	5-Bromo-dideoxyuridine	353.1	5-Br-ddU:dA	666.3
C	5-Methyl-dideoxycytidine	287.2	5-Me-ddC: 7-deaza-dG	615.4
A	Dideoxyadenosine	297.2	ddA:dT	601.4
G	Dideoxyinosine	298.2	ddl:dC	587.4

The positional mass difference at every base position is given by:

$$(xii) \quad M_{diff}[n] = M_{pair}[n] - M_{lightest},$$

where $M_{diff}[n]$ is the n^{th} positional mass difference, $M_{pair}[n]$ is the mass of n^{th} terminating base pair, and $M_{lightest}$ is the mass of the lightest terminating base pair in daltons. Substituting in the values from the table above yields:

$$M_{diff}["G"] = (587.4 - 587.4) = 0$$

$$M_{diff}["A"] = (601.4 - 587.4) = 14$$

$$M_{diff}["C"] = (615.4 - 587.4) = 28$$

$$M_{diff}["T"] = (666.3 - 587.4) = 78.9$$

Since each terminating base pair has a unique positional mass difference, the base sequence can be determined unambiguously. The maximum mass shift in this case is 78.9 daltons. When choosing a set of terminating nucleotides it is

-44-

important to select the set such that the positional mass difference for each base termination is distinct and resolvable by mass.

If modified nucleotide terminators are not used, it is still possible to implement Forced Mass Modulation by carrying out each of the four termination reactions separately using mass-labeled primers rather than modified terminators, combining all reaction products, and then obtaining a mass spectrum. In order to produce the same positional mass differences as shown in Example 1, using a set of pair-matched nucleotides and the standard dideoxy terminators, the following primer mass shifts are required:

10	<u>Termination Reaction</u>	<u>Primer Mass</u>
	C	"reference" primer
	T	reference primer + 15 Da
	A	reference primer + 24 Da
	G	reference primer + 39 Da

15 This method is essentially equivalent to multiplexing four single-nucleotide sequencing reactions in the same spectrum, except that all the sequencing products originate from the same priming site but terminate on different nucleotides.

EXAMPLE 3

20 **Forced Mass Modulation in the Detection and Scoring of Single Nucleotide Polymorphisms**

Forced Mass Modulation can be used to simplify the analysis of closely related sequence variants, as is required in the detection and scoring of single nucleotide polymorphisms. Figure 6 shows three sequence variants that differ from each other only at a single base position sequenced by a conventional Sanger reaction. The mass distribution of the reaction products is so complex that it can be uninterpretable, even if the base sequences of the variants are known *a priori*.

Figure 7 shows the same three variants sequenced by Forced Mass Modulation using mass-matched deoxynucleotides (dN = 310 Da) and the standard dideoxy terminators. The positions and identities of the single-nucleotide changes are immediately apparent from the mass spectrum.

-45-

Since the masses of the sequencing reaction products are constrained to fall within the shaded regions of the spectrum in Figure 7b, it is possible to multiplex other sequences on the same spectrum.

EXAMPLE 4

5 Base Composition Density Distributions for the Total Set of possible 7-base Oligonucleotides

For this implementation, three sets of 7-base oligonucleotides comprising all possible base compositions for a 7-base oligonucleotide can be obtained; the first set comprising the four natural bases (dA, dG, dC and dT), the second set
10 comprising three of the natural bases (dA, dC and dT) and the nucleotide analog 7-deaza-deoxyguanosine (7-deaza-dG) substituted for dG, and the third set comprising three of the natural bases (dA, dG and dC) and the nucleotide analog deuterio-deoxythymine (deutero-dT) substituted for dT. Figure 8 shows the actual base composition density distributions for the total set of possible 7-base
15 oligonucleotides using the three different nucleotide sets. Note that for the set of naturally occurring bases (Figure 8a), nearly every base composition has its own distinct mass value, but most of these mass values are spaced only one dalton from each other. Increasing the peak separation to three daltons by substitution of dG with 7-deaza-dG (Figure 8b) markedly increases the average
20 number of base compositions per observed mass, particularly for those masses in the center of the range, but any two oligonucleotides of the same length with different molecular weights will have to be separated by at least three daltons. Similarly, substitution of dT with deuterio-dT (Figure 8c) gives a minimum peak separation between oligonucleotides having the same length but different
25 molecular weights of eight daltons. The trade-off for a greater peak separation is a greater number of oligonucleotides that have exactly the same mass for a given oligonucleotide length.

Since modifications will be apparent to those of skill in this art, it is
30 intended that this invention be limited only by the scope of the appended claims.

-46-

WHAT IS CLAIMED IS:

1. A method for identifying the nucleotide at one or more base positions in a target nucleic acid molecule, comprising:
 - synthesizing extension products of the target nucleic acid in the presence
 - 5 of chain terminating nucleotides and mass-matched nucleotides;
 - determining the mass of each extension product; and
 - calculating a mass shift from a period for the mass of each extension product,
 - whereby nucleotide(s) at one or more base positions is determined by
 - 10 identifying the nucleotide that corresponds to each mass shift.
2. The method of claim 1 that is a method for determining a nucleotide sequence of a target nucleic acid, comprising:
 - synthesizing extension products of the target nucleic acid in the presence
 - of chain terminating nucleotides and mass-matched nucleotides;
 - 15 determining the mass of each extension product; and
 - calculating a mass shift from a period for the mass of each extension product,
 - whereby the nucleotide sequence of the target nucleic acid is determined by assigning a nucleotide corresponding to each mass shift.
 - 20
3. The method of claim 1, wherein the mass-matched deoxynucleotides are identical.
4. The method of claim 1, wherein a mass-matched deoxynucleotide is deoxyinosine, 5-nitroindole, 3-nitropyrrole, 3-methyl 7-propynyl isocarbostyryl, 5-methyl iscarbostyryl or 3-methyl iscarbostyryl.
- 25 5. A method for identifying nucleotides at one or more base positions in a plurality of target nucleic acids molecules, comprising:
 - synthesizing extension products of the target nucleic acid in the presence
 - of chain terminating nucleotides and mass-matched nucleotides;
 - determining the mass of each extension product; and
 - 30 calculating a mass shift from a period for the mass of each extension product,

-47-

whereby the nucleotides in the target nucleic acid molecules are identified by determining the nucleotide that corresponds to each mass shift.

6. The method of claim 5 that is a method for determining nucleotide sequences of a plurality of target nucleic acids molecules, comprising:

- 5 synthesizing extension products of the target nucleic acid in the presence of chain terminating nucleotides and mass-matched nucleotides;
determining the mass of each extension product; and
calculating a mass shift from a period for the mass of each extension product,

- 10 whereby the nucleotide sequences of the target nucleic acids are determined by determining the nucleotide that corresponds to each mass shift.

7. The method of claim 5, wherein the mass-matched deoxynucleotides are identical to one another.

8. The method of claim 1, wherein a mass-matched deoxynucleotide
15 is deoxyinosine, 5-nitroindole, 3-nitropyrrole, 3-methyl 7-propynyl iscarbostyryl, 5-methyl iscarbostyryl or 3-methyl iscarbostyryl.

9. A method for identifying nucleotides at one or more base positions in a plurality of target nucleic acids molecules, comprising:

- 20 synthesizing extension products of the target nucleic acid in the presence of chain terminating nucleotides and mass-matched nucleotides;
determining the mass of each extension product; and
calculating a mass shift from a period for the mass of each extension product,

- 25 whereby the nucleotides in the target nucleic acid molecules are identified by determining the nucleotide that corresponds to each mass shift.

10. A method for determining a nucleotide sequence of a target nucleic acid molecule, comprising:

- incorporating pair-matched nucleotides into the target nucleic acid;
synthesizing extension products of the target nucleic acid in the
30 presence of a partially duplex hairpin primer, chain terminating nucleotides and pair-matched nucleotides;
determining the mass of each extension product; and

-48-

calculating a mass shift from a period for the mass of each extension product;

whereby the nucleotide sequence of the target nucleic acid is determined by assigning a nucleotide corresponding to each mass shift.

5 11. The method of claim 10, wherein the chain terminating nucleotides are mass-matched.

12. The method of claim 10, wherein the chain terminating nucleotide base pairs have distinct molecular weights.

13. A method for determining nucleotide sequences of a plurality of
10 target nucleic acids, comprising:

incorporating pair-matched nucleotides into the target nucleic acids;

synthesizing extension products of the target nucleic acids in the presence of a partially duplex hairpin primer, chain terminating nucleotides and
15 pair-matched nucleotides;

amplifying the target nucleic acid sequences in the presence of pair-matched nucleotides;

determining the mass of each extension product; and

calculating a mass shift from a period for the mass of each
20 extension product;

whereby the nucleotide sequences of the target nucleic acids are determined by assigning a nucleotide corresponding to each mass shift.

14. The method of claim 13, wherein the chain terminating nucleotides are mass-matched.

25 15. The method of claim 13, wherein the chain terminating nucleotide base pairs have distinct molecular weights.

16. The method of claim 13, wherein the primers are mass-labeled.

17. A method for detecting a one or a plurality of target nucleic acid(s) or one or plurality of nucleotides therein molecules, comprising:

30 (a) copying the target nucleic acid molecule(s) in the presence of a pair-matched set of nucleotides;

-49-

(b) denaturing the resulting copies of the target(s) to produce single-stranded templates;

(c) annealing and ligating one or a plurality of partially duplex hairpin primers to the single-stranded template(s);

5 (d) extending the primer(s) in the presence of chain terminating nucleotides and pair-matched nucleotides to produce extension products, wherein the extension products follow a periodic mass distribution that is determined by the mass of the pair-matched nucleotide set; and

10 (e) detecting each of the targets or nucleotides therein by virtue of the mass shift of each extension product from its corresponding periodic reference mass.

18. The method of claim 17, wherein the chain terminating nucleotides are mass-matched.

15 19. The method of claim 17, wherein the chain terminating nucleotide base pairs have distinct molecular weights.

20. The method of claim 17, wherein the primers are mass-labeled.

21. A kit for determining the sequence of a target nucleic acid, comprising mass-matched nucleotides.

20 22. A kit for determining the sequence of a target nucleic acid, comprising pair-matched nucleotides and mass-matched chain terminating nucleotides.

25 23. A kit for determining the sequence of a target nucleic acid, comprising pair-matched nucleotides and chain terminating nucleotides that form base pairs of distinct molecular weight, and optionally including instructions for sequencing using these reagents.

24. A kit for determining the sequence of a target nucleic acid, comprising pair-matched nucleotides and mass-labeled primers, and optionally including instructions for sequencing using these reagents

30 25. A method for detecting different nucleotide base compositions in a population of nucleic acids having identical length, comprising:

synthesizing the nucleic acids in the presence of one or more nucleotide analogs to produce synthesized nucleic acids; and

-50-

- determining a mass of each synthesized nucleic acid;
whereby different nucleotide base compositions are detected by
determining the mass of each synthesized nucleic acid,
wherein the nucleotide analog separates the masses of nucleic acids
5 having different base compositions in a predetermined interval.
26. The method of claim 25, wherein the population of nucleic acids
having identical length and different base compositions differ in base
composition by a single base.
27. A method for detecting a plurality of target nucleic acid molecules
10 in a sample containing nucleic acid molecules, comprising:
preparing a composition containing plurality of pair-matched nucleic acid
molecules or mass-matched nucleic acid molecules from a sample comprising the
target nucleic acid molecules;
analyzing the resulting composition by mass spectrometry; and
15 detecting target nucleic acid molecules.
28. A process for detecting a mutation in a target nucleic acid sequence
in a target nucleic acid molecule, in a sample, comprising:
a) hybridizing a nucleic acid molecule a primer to nucleic acid
molecules in the sample, thereby producing a hybridized primer wherein:
20 the nucleic molecules from the sample are optionally immobilized;
the primer is complementary to a sequence in the target nucleic acid
sequence that is adjacent to the region suspected of containing a mutation
sequence;
- b) contacting the hybridized primer with a composition comprising
25 mass-matched deoxyribonucleoside triphosphates and a chain terminating
nucleotide selected from a dideoxyribonucleoside triphosphate or a
3'-deoxynucleoside triphosphate and optionally one or more
deoxyribonucleoside triphosphates, such that the hybridized primer is
extended until a chain terminating nucleotide is incorporated, thereby
30 producing an extended primer; and

-51-

c) determining the mass of the extended primer, thereby determining whether a mutation is present in the target nucleic acid sequence.

29. The process of claim 28, wherein the chain terminating
5 nucleotides are mass-matched.

30. The method of claim 28, wherein the mass of the extended primer is determined by mass spectrometry.

31. A process for detecting mutations in a plurality of target nucleic acid sequences in a sample, comprising:

10 a) hybridizing a plurality of primers to nucleic acid molecules in the sample, thereby producing a hybridized primers, wherein: the nucleic molecules from the sample are optionally immobilized; each primer is complementary to a sequence of a target nucleic acid sequence that is adjacent to a region suspected of containing a mutation
15 sequence;

b) contacting the hybridized primers with a composition comprising a chain terminating nucleotide selected from a mass-matched dideoxynucleoside triphosphate or a 3'-deoxynucleoside triphosphate and one or more deoxyribonucleoside triphosphates, such that the
20 hybridized primers are extended until a chain terminating nucleotide is incorporated, thereby producing an extended primer; and

c) determining the mass of the extended primers, thereby determining whether mutations are present in the target nucleic acid sequences.

25 32. The process of claim 31, wherein the chain terminating nucleotides are mass-matched.

33. The method of claim 31, wherein the mass of the extended primers are determined by mass spectrometry.

34. A method for detecting a target nucleic acid sequence, comprising
30 the steps of:

a) hybridizing a primer to a nucleic acid molecule comprising a target nucleic acid sequence, wherein the primer can be extended in a 3'

-52-

direction towards the target nucleic acid sequence, and wherein the 5' end of the primer can be selectively cleaved from the extension product;

b) extending the primer in the presence of mass matched deoxyribonucleotides and a polymerase to produce an extension product;

5 c) selectively cleaving the 5' end of the primer from the extension product to produce a portion of the primer and a cleaved extension product; and

d) detecting the cleaved extension product.

35. The method of claim 34, wherein the cleaved extension product is
10 detected by mass spectrometry.

36. A method for detecting a plurality target nucleic acid sequence, comprising the steps of:

a) hybridizing a primer or plurality thereof nucleic acid molecules comprising target nucleic acid sequences, wherein the primers can be
15 extended in a 3' direction towards the target nucleic acid sequence, and wherein the 5' end of the hybridized mass-matched nucleic acid molecules can be selectively cleaved from the extension product;

b) extending the primers in the presence of mass matched deoxyribonucleotides and a polymerase to produce extension products;

20 c) selectively cleaving the 5' end of the primers from the extension products to produce portions of the primers and cleaved extension products; and

d) detecting the cleaved extension products.

37. The method of claim 36, wherein the cleaved extension product is
25 detected by mass spectrometry.

38. A method for detecting a target nucleic acid sequence, comprising the steps of:

a) hybridizing to a nucleic acid molecule comprising the target nucleic acid sequence

30 a first primer, which can be extended in a 3' direction towards the target nucleic acid sequence, and wherein the 5' end

-53-

of the primer can be selectively cleaved from the extension product, and

a second primer, which can be extended in a 3' direction towards the first primer;

5 b) extending the primers in the presence of mass-matched nucleotides to produce a double stranded amplification product;

c) selectively cleaving the 5' end of the first primer in the amplification product, to produce a double stranded amplification product comprising a cleaved primer extension product comprising a 5' portion
10 and a 3' portion;

d) denaturing the product of step c); and

e) detecting the 3' portion of the cleaved primer extension product.

39. The method of claim 38, wherein the cleaved extension product is
15 detected by mass spectrometry.

40. A method for detecting a plurality target nucleic acid sequences, comprising:

a) hybridizing to each of a plurality of nucleic acid molecules comprising the target nucleic acid sequence

20 a first primer, which can be extended in a 3' direction towards the target nucleic acid sequence, and wherein the 5' end of the primer can be selectively cleaved from the extension product, and

25 a second primer, which can be extended in a 3' direction towards the first primer;

b) extending the primers in the presence of mass-matched nucleotides or pair-matched nucleotides to produce double stranded amplification products;

30 c) selectively cleaving the 5' end of each of the first primers in the amplification product, to produce double stranded amplification products comprising cleaved primer extension products comprising a 5' portion and a 3' portion;

-54-

- d) denaturing the products of step c); and
- e) detecting the 3' portions of the cleaved primer extension products by virtue of the masses.

41. The method of claim 40, wherein detection is effected by mass spectrometry.

42. A method for detecting a target nucleic acid sequence, comprising

- a) hybridizing first and second primers to a nucleic acid molecule containing the target nucleic acid sequence, wherein a primer contains a selectively cleavable site at its 3' end;
- b) extending the primers in the presence of mass-matched nucleotides;
- c) cleaving the resulting product at the selectively cleavable sites;
- d) analyzing the masses of the cleavage products, whereby the target sequence is detected.

43. The method of claim 42, wherein the cleaved extension product is detected by mass spectrometry.

44. The process of claim 43, wherein a plurality of primers are hybridized and a plurality of target sequences are identified in a single reaction.

45. The method of claim 44, wherein the cleaved extension products are detected by mass spectrometry.

46. A computer-based method for identifying nucleotide or nucleotides at one or more base positions in a target nucleic acid molecule or plurality thereof, comprising:

- a) entering the primer sequence or primer mass, the mass of an individual mass-matched deoxynucleotide into the computer and the identify of chain terminators used;
- b) entering the masses of the fragments generated by a primer extension reaction, wherein the primer is extended by mass-matched deoxynucleotides;
- c) determining P_{base} , wherein P_{base} is the base periodicity in daltons;
- d) calculating $M_{diff}[n]$ for each nucleotide base to be identified, wherein:

$$M_{diff}[n] = M_{obs}[n] - M_{pr}[n];$$

-55-

$$M_{PR}[n] = (M_{primer} + M_{light}) + (n - 1) P_{base};$$

$M_{obs}[n]$ is the observed peak;

where:

n is the base position;

5 $M_{PR}[n]$ is the n^{th} periodic reference mass;

M_{primer} is the mass of the primer;

M_{light} is the mass of the lightest nucleotide terminator; and

e) determining the identity of a nucleotide at any base position or the positional mass difference by determining $M_{diff}[n]$ and comparing it to a database
10 of previously calculated values of M_{diff} for each of the chain terminating nucleotides.

47. A system for high throughput analysis of nucleic acid samples, comprising:

a processing stations that performs a chain extension reaction, in the
15 presence of mass-matched nucleic nucleotides, on a nucleic acid sample in a reaction mixture;

a robotic system that transports the resulting products from the processing station to a mass measuring station, wherein the masses of the products of the reaction are determined; and


20 a data analysis system that processes the data from the mass measuring station by performing the method of claim 46 to identify a nucleotide or nucleotides at one or more base positions in nucleic acid molecule in the sample.

48. The system of claim 47, further comprising a control system that determines when processing at each station is complete and, in response, moves
25 the sample to the next test station, and continuously processes samples one after another until the control system receives a stop instruction.




49. The system of claim 46, wherein the mass measuring station is a mass spectrometer.

a) Sequence #1: AGCGTA

Primer	Extension	Mass (Da)
3016 Da	Products	
—	agcgta	4878.2
—	agcgt	4565.0
—	agcg	4260.8**
—	agc	3931.6
—	ag	3642.4*
—	a	3313.2

b) Sequence #2: GATCCT

Primer	Extension	Mass (Da)
3016 Da	Products	
—	gagcct	4854.2
—	gagcc	4550.0
—	gagc	4260.8**
—	gag	3971.6
—	ga	3642.4*
—	g	3329.2



e)

```

gcgct
GCGTA
ggcct
ggcta
acgct
acgta
AGCCT
agcta


```

Figure 1

a) Nucleotide Mass (Da) Sequence #1: GCATG

ddN	310
ddC	273
ddT	288
ddA	297
ddG	313

Primer #1	Extension Products	Mass (Da)
3327 Da	nnnng	4880
	nnnt	4545
	nna	4244
	nc	3910
	g	3640

b) Sequence #2: CATGC

Primer #2	Extension Products	Mass (Da)
3404 Da	nnnnc	4917
	nnng	4647
	nnt	4312
	na	4011
	c	3677




Figure 2

Sequence #1: GCATA

Primer #1 3327 Da	Extension Products	Mass (Da)
————	nnnna	4864
————	nnnt	4545
————	nna	4244
————	nc	3910
————	g	3640

Sequence #3: CATGC

Primer #3 3404 Da	Extension Products	Mass (Da)
————	nnnnc	4917
————	nnng	4647
————	nnt	4312
————	na	4011
————	c	3677

Sequence #2: TCAGG

Primer #2 3481 Da	Extension Products	Mass (Da)
————	nnnng	5034
————	nnng	4724
————	nna	4398
————	nc	4064
————	t	3769

Sequence #4: AACTC

Primer #4 3558 Da	Extension Products	Mass (Da)
————	nnnnc	5071
————	nnnt	4776
————	nnc	4451
————	na	4165
————	a	3855


Sequence

Figure 3




Figure 4




Figure 5




Figure 6





Figure 7

Figure 8


Base composition density distributions for
7-mers using different nucleotide sets.



C = 289.2
T = 304.2
A = 313.2
G = 329.2

Naturally Occurring Bases


Peaks can be closer than one dalton.
Total No. of different base compositions = 120
Actual number of represented masses = 110
Avg. No. of compositions per mass value = 1.091



C = 289.2
T = 304.2
A = 313.2
G = 328.2

Substitution with 7-deaza-dG

Minimum peak separation = 3 daltons
Number of allowed mass values = 92
Actual number of represented masses = 64
Avg. No. of compositions per mass value = 1.875



C = 289.2
T = 305.2
A = 313.2
G = 329.2

Substitution with deuterio-dT

Minimum peak separation = 8 daltons
Number of allowed mass values = 36
Actual number of represented masses = 34
Avg. No. of compositions per mass value = 3.529

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.